

Study of Distribution of ABO and Rh Blood Groups among First Year Medical Students in CAIMS Medical College, Karimnagar

D. Lavanya¹, A. R. Vidyullatha^{2*}, Akshay Berad³

¹Assistant Professor, Dept. of Physiology, Sri Venkateswara Medical College, Tirupati, AP, India

²Assistant Professor, Dept. of Physiology, CAIMS, Karimnagar, Telangana state, India

³Associate Professor, Dept. of physiology, RIMS, Adilabad, Telangana state, India

Original Research Article

*Corresponding author

A. R. Vidyullatha

Article History

Received: 11.11.2018

Accepted: 20.11.2018

Published: 30.11.2018



Abstract: An important milestone in the history of blood transfusion was the discovery of the ABO blood groups by Karl Landsteiner, followed by discovery of Rh (D) antigen. Landsteiner's achievements lead to several discoveries in the field of immunohaematology. The ABO blood group is useful in selecting the appropriately matched blood group during blood transfusion, organ transplantation, finding out association of blood groups and diseases etc. ABO blood group system classifies blood groups of people into four different types namely A, B, O and AB. The need for the study of frequency distribution of blood group is multipurpose. This study was done to know the frequency distribution of blood groups among first year medical students. This study was conducted on 100 medical students in the Department of Physiology at CAIMS medical college in Karimnagar Telangana state. The blood samples were collected by finger prick method. ABO blood grouping and Rhesus factors (Rh) typing determined by glass slide method. Out of total 100 medical students who volunteered, 30 were males and 70 females. In our study, the most frequently occurring blood group was O (42%) followed by B group (29%), A group (17%) and AB group (12%). 81% students were Rh positive whereas only 19% were Rh negative. This study may enable us to contact individuals belonging to a particular blood group at times of medical emergencies when blood transfusion is required.

Keywords: ABO, Rh-D, Blood group system.

INTRODUCTION

ABO and Rhesus (Rh) blood group systems till today remain clinically most important inspite of being identification of around twenty nine blood group systems, enumerated by International Society of Blood Transfusion. In 1900, Karl Landsteiner detected the human ABO blood group [1]. The Rh blood group system was discovered during 1939–1940 by Landsteiner, Weiner, Levine and Stetson, clarifying the basis of many unpredicted transfusion reactions. In 1945, Coombs, Mourant and Race described the use of antihuman globulin (Coombs test) for incomplete antibodies [2]. Later, these two systems have substantiated to be the most important in transfusion medicine. Today, the requirement for blood group frequency and prevalence studies is multiuse, besides their importance in relative to blood transfusion and organ transplantation. Human red blood cells contain on their surface a series of glycoproteins and glycolipids, which constitute blood group antigens. Development of these antigens are genetically controlled and they appear early in fetal life and remain unchanged till death [3]. The major ABO blood group system is divided into four blood types on the basis of presence or

absence of A and B surface antigens. The blood groups are A, B, O and AB. The frequency of four main ABO blood groups varies in the population throughout the world. ABO blood group system derives its importance from the fact that A and B are strongly antigenic and anti A and anti B naturally occurring antibodies present in the serum of persons lacking the corresponding antigen and these antibodies are capable of producing intravascular hemolysis in case of incompatible transfusion [4]. Rh antigens are highly immunogenic and till now 49 Rh antigens are identified. D antigen is most significant and D negative individuals produce anti-D if they encounter the D antigen through transfusion or pregnancy and causes hemolytic transfusion reaction, or hemolytic disease of fetus and newborn. For this reason, the Rh status is routinely determined in blood donors, transfusion recipients, and in mothers-to-be [5]. Acquiring knowledge on distribution of ABO blood groups at local and regional levels will not only be helpful in the effective management of blood banks but also in safe blood transfusion services [6]. Hence, the present study was planned with the aim of determining the distribution of ABO and Rh blood groups in healthy young adults in

college. This study may enable us to contact individuals belonging to a particular blood group at times of medical emergencies when blood transfusion is required.

MATERIALS AND METHODS

Study was conducted on first year MBBS medical students in the Department of Physiology at CAIMS Medical College in Karimnagar Telangana State. Total of 100 medical students, 30 males and 70 females, samples were collected by finger prick method under aseptic precautions. The ABO blood grouping and Rhesus factors (Rh) typing determined by glass slide method, which is based on antigen antibody agglutination. Commercially available standard anti sera A, anti sera B and anti sera D were used for the study. RBCs suspended in isotonic saline were treated with anti-A, anti-B and anti-D anti sera on separate glass slides, marked as A, B and D and mixed with separate applicator sticks. The mixture observed for agglutination, both macroscopically and microscopically and compared with the control. The

blood group was determined based on agglutination with the corresponding anti sera. If agglutination was present in the blood drop A, then it belongs to A blood group, agglutination in blood drop B, B group, agglutination in both A and B blood drops, AB group and if no agglutination in both A and B drops, then O group. Similarly, agglutination in blood drop D was considered as Rh Positive and no agglutination Rh negative. The data was expressed as percentages.

RESULTS

Out of total 100 participants 70 were female students and 30 were male students (Table-1).

Table-2 Shows that most common blood group was O (42%) followed by B (29%), A (17%), AB (12%).

Table-3 shows that 81 % students were Rh positive and 19 % were Rh negative.

Table-1: Gender and number of participant students

Gender	Number of students (100)
Male	30
Female	70

Table-2: Distribution of ABO blood group among students

Blood groups	Number of students	Percentage of students
A	17	17%
AB	12	12%
B	29	29%
O	42	42%

Table-3: Distribution of Rh blood group among students

Blood Group	Number of students	Percentage
Rh Positive	81	81%
Rh Negative	19	19%

Table-4: Details of ABO and Rh blood group distribution among 100 students

Blood groups	Number of students (100)	Percentage
A Rh Positive	15	15%
B Rh Positive	25	25%
AB Rh Positive	10	10%
O Rh Positive	31	31%
A Rh Negative	2	2%
B Rh Negative	4	4%
AB Rh Negative	2	2%
O Rh Negative	11	11%

Table-4 shows O + is (31%) commonest blood group among 100 students and A- and AB – are rarest blood groups (2%).

DISCUSSION

Blood grouping is an important parameter for social, professional and medical needs. In the present study, the frequency of blood group O was the highest (42%) and the least frequency was that of blood group AB (12 %). Several studies within the Telangana state [7] and rest of India [8-12] and other countries [13-15] have reported variations in the blood group distribution in diverse populations. This study helped in analyzing the distribution of blood groups among college students who comprise a heterogeneous population within this region.

CONCLUSION

The study confirms that blood group O was the commonest of the ABO blood group system among the medical students studied and AB blood group was the least. Rhesus positive was commoner than Rhesus negative. Knowledge of blood group distribution is important for clinical studies, for reliable geographical information, blood bank management and for forensic studies in the population. Such a study would create awareness about self-blood grouping and also enable one to prepare a database of the available blood groups which can be utilized during medical emergencies for safe blood transfusion.

REFERENCES

1. Garratty, G., Dzik, W., Issitt, P. D., Lublin, D. M., Reid, M. E., & Zelinski, T. (2000). Terminology for blood group antigens and genes—historical origins and guidelines in the new millennium. *Transfusion*, 40(4), 477-489.
2. Hillyer, C. D. (2013). Blood banking and transfusion medicine—history, industry, and discipline. In *Transfusion Medicine and Hemostasis (Second Edition)* (pp. 3-9).
3. Firkin, F., Chesterman, C., Penington, D., & Rush, B. (1989). Blood groups; blood transfusion; acquired immune deficiency syndrome. *De Gruchy's Clinical Hematology in Medical Practice*, 5, 475-96.
4. Harmening, D. M., & Firestone, D. (2012). The ABO blood group system. *Modern blood banking and transfusion practices*, 5, 108-32.
5. Dean, L. (2005). Blood Groups and Red Cell Antigens, a guide to the differences in our blood types that complicate blood transfusions and pregnancy. *National Center for Biotechnology Information*.
6. Patel, P. (2011). Frequency and distribution of blood groups in blood donors in western Ahmedabad—a hospital based study. *blood*, 202-206.
7. Vujhini, S. K., Sanagapati, P. R. R., & Kandukuri, M. (2014). Frequency and distribution of blood groups in blood donors of a tertiary care hospital in Rangareddy district of Telanaga State, India.
8. Rao, N. M., Grandhi, B., Shanthi, V., Byna, S. S. R., Sreesailam, S., & Murthy, B. K. (2015). Study of Blood Groups in Blood Donors in Narayana Medical College & Hospital, Nellore (AP), India. *International Journal of Current Medical And Applied Sciences*, 6(2), 110-113.
9. Krishna, M. C., Sharadha, M. S., Hulinaykar, R. M., & Harish, S. G. (2014). Frequency and distribution of ABO and Rhesus (D) blood groups in and around Tumkur, Karnataka, Study from Teritiary Care Teaching Hospital. *Internat Jour. Healthcare Sci*, 2(1), 135-9.
10. Purandare, V. R., & Prasad, N. B. (2012). Distribution of ABO blood groups in healthy young adults in Pune city. *Int. J. Basic Appl. Med. Sci*, 2(3), 74-78.
11. Panmei, K., Yumnam, P., & Ngaomei, G. (2014). Frequency Distribution of ABO and Rh blood groups among students of Maram tribe of Don Bosco College, Maram, Manipur. *Int J Pure App Biosci*, 2(4), 61-6.
12. Sidhu, S. (2003). Distribution of the ABO blood groups and Rh (D) factor among the scheduled caste population of Punjab. *The Anthropologist*, 5(3), 203-204.
13. Bamidele, O., Arokoyo, D. S., & Akinbola, A. O. (2013). Distribution of ABO and rhesus blood groups among medical students in Bowen University, Iwo, Nigeria. *Ann. Biol. Res*, 4(11), 1-6.
14. Sultana, R., Rahman, Z., Helali, A. M., Yousuf, R., Mustafa, S., Salam, A., & Haque, M. (2013). Study of ABO and RH-D Blood Groups among the common people of capital city of Bangladesh. *International Journal of Pharmacy and Pharmaceutical Sciences*, 5(3), 814-816.
15. Hammed, A., Hussain, W., Ahmed, J., Rabbi, F., & Quersh, J. A. (2002). Prevalence of phenotypes and Genes of ABO and Rhesus (Rh) blood groups in Faisalabad, Pakistan. *Pak J Biol Sci*, 5(6), 722-724.