∂ OPEN ACCESS Scholars International Journal of Obstetrics and Gynecology

Abbreviated Key Title: Sch Int J Obstet Gynec ISSN 2616-8235 (Print) |ISSN 2617-3492 (Online) Scholars Middle East Publishers, Dubai, United Arab Emirates Journal homepage: https://saudijournals.com/journal/sijog/home

Original Research Article

Female Urinary Incontinence – Epidemiology, Causes, Risk Factors and Management in Tribal Rural Women of South Rajasthan

Dr. Koshalya Bordia¹, BRIG Dr. PK Bhatnagar²*

¹Professor, Department of Obstetrics & Gynecology, Pacific Institute Of Medical Sciences (PIMS), Ambua Road, RJ SH 32, Village-Umarda, Udaipur, Rajasthan 313015, India

²Venue of Research, Pacific Institute of Medical Sciences, Ambua Road, RJ SH 32, Village-Umarda, Udaipur, Rajasthan 313015, India

DOI: 10.36348/sijog.2019.v02i07.001

| Received: 29.06.2019 | Accepted: 08.07.2019 | Published: 15.07.2019

*Corresponding author: Dr. PK Bhatnagar

Abstract

Urinary incontinence has great bearing on quality of life, social life and is of high public health importance [1]. 12 % women of all age and 27% of the women above 50 year of age suffer from urinary incontinence [2]. the risk factors have been history of gynae operations, prolonged labor difficult labor, forceps or vacuum delivery, repeated deliveries, chronic cough ,chronic constipation and failure to get early treatment [3]. Aim of present study has been to determine role of risk factors and to suggest preventive methods, early diagnosis and treatment.

Keywords: Urinary Incontinence, Stress Urinary Incontinence, Urge Incontinence, Urinary Fistulae Prolapse Uterus, Rural women Health.

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (Non-Commercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

The International Continence Society defined Urinary incontinence as "the complaint of any involuntary leakage of urine and which is a social or hygienic problem [4] as an estimate 400 million people around the world are sufferers and it increases with age [5]. 7% of women 20–40 years of age, 17% of 40–60 years of age, 23% of 60–80 years of age have been found to have moderately severe urinary incontinence [6] risk factors for Urinary incontinence include increasing age, parity, vaginal deliveries, obesity, surgery, constipation, and chronic respiratory problems such as cough [7].

The inability to control urine is quite an unpleasant and distressing problem. Although it does not lead to death, it causes substantial morbidity, social seclusion, and psychological stress resulting in impaired Quality of life. Many women are too embarrassed to talk about it and some believe it to be untreatable [8]. Routine speculum examination and simple tests like Qtip test, pad test and ultrasonography emphasizing on post residual volume can help to know the type of incontinence and factors leading to incontinence [9] conservatively Pelvic Floor Muscle Training remains the first-line treatment for female urinary incontinence [10]. The modified intravaginal slingplasty is a new and interesting procedure [11]. It has replaced the earlier kellys sutures and anterior colporrhaphy and traditional abdominal passage of a needle or packing forceps, to retrieve the sling and tie at proximal urethra [12].

MATERIAL AND METHODS

This is a prospective observational study done in a Tertiary Care Center, hospital of PIMS (PACIFIC INSTITUTE OF MEDICAL SCIENCES, VILLAGE UMARDA, UDAIPUR, RAJASTHAN from 2015 to 2019 Women who attended gynae opd with urinary complains of incontinence were interviewed. Risk factors were noted. Age, parity, socioeconomic status, education, age of marriage and residential address were noted. Complains and details of illness, obstetric history, medical and surgical history was noted. General examination included presence of anemia, blood pressure and edema. Systemic examination included cardiac and lung condition. Diagnostic urodyanamic study, uroflowmetery standing stress test resting and stress urethral pressure profile and medium filled cystometery. Ultrasonic examination results were recorded. Blood HB, SUGAR, UREA, CREATININE, BLOOD COUNTS, BT, CT, PT, HIV, HBSAG, VDRL, URINE SUGAR and ALBUMIN, BLOOD GROUPING results recorded. RISK FACTORS AND OUTCOME of conservative and operative treatment was recorded.

OBSERVATIONS

5. INO	Age in years	No of patients	Percentage
1	30 and less	0	0
2	30-60	18	40
3	61-80	27	60
4	Parity 2 and below	8	17.77
5	Parity 3 and above	37	82.22
6	High socio economic	0	
7	Low socio economic	45	100
8	Education below 5	45	100
9	Education above 6	0	

 S
 No
 A go in woorg
 No of potients
 Bereastage

Table-2: Clinical	presentation and	l risk factors

S. No	Types Of Incontinence And Causes	Number	Percentage
1	Stress urinary incontinence	28	62.22
2	mixed urinary incontinence	7	15.55
3	Urge incontinence	10	22.22
4	History of gyne surgery	5	11.11
5	Forceps vacuum prolonged labor	15	33.33
6	Diabetic with repeated urinary inf	5	11.11
7	Chronic cough	12	26.66
8	constipation	8	17.77
9	Relief with conservative	5	11.11
10	Relief with vaginal repair	15	33.33
11	Relief with sling surgery	25	55.55

RESULTS

Total 175 patients were studied and 45 were detected to have urinary incontinence (25.71%)18(40%) patients were 30-60 years of age,27 (60%) were 61-80 years of age.8(17.77) were para 2 or below, 37(82.22%) were of 3 and more parity. All patients were from low socioeconomic status and were illiterate. 28 (62.22%) had stress urinary incontinence, 7(15.55%) were mixed and 10(22.22%) had urge incontinence. There was history of gynae surgery in 5(11.11%), 15(33.33%) had prolonged labor, forceps or vacuum delivery.5 (11.11%) were diabetic with repeated urinary infections.12 (26.66%) had chronic cough, 8(17.77%) had constipation. 5(11.11%) were successfully treated with pelvic floor muscle training, 15(33.33%) were treated by definitive surgery and vaginal repair, 25(55.55%) got relief from sling surgery.

DISCUSSION

In the present study 45 (25.71%) patients were found to have urinary incontinence. 30% incidence was found by Herzog *et al.*, [13]. Brown *et al.*, [14] found a much higher incidence (55%) and Black *et al.*, [15] higher still (62%).In our study stress incontinence 28(62.22%), mixed 7(15.55%) and urge 10(22.2%) was reported. 13% reported by Brown *et al.*, In our study incontinence was common amongst menopausal women viz, 37(82.22%) whereas Fultz *et al.*, [16] reported a 30-40% incidence. In our study average age was 45 years and mean parity 4 whereas in a study conducted by Swift and Ostergard [17], mean age and mean parity were 59.8 and 2.3 respectively. Prolonged repeated difficult labor was detected as main risk factor 15(33.33%). Others included vaginal hysterectomy, colporrhaphy (11.11%), chronic cough 12(17.77%) and 5(11.11%) had diabetes and repeated lower urinary tract infections. Majority got relieved 25(55.55%) and 15(33, 33%) were benefited by vaginal repair.

CONCLUSION

Prevalence of Urinary Incontinence is high in rural women. Stress Urinary Incontinence is more common. Generating awareness regarding UI may help to improve health-seeking behavior. There is significant suffering and social life changes in women with Urinary incontinence. History and speculum examination is diagnostic. Conservative pelvic floor muscle training is preventive and sling operation is curative .Institutional deliveries, good antenatal care and preventive pelvic floor muscle training can help reducing incidence and suffering of rural women.

REFERENCES

- Abrams, P., Andersson, K. E., Birder, L., Brubaker, L., Cardozo, L., Chapple, C., ... & Drake, M. (2010). Fourth International Consultation on Incontinence Recommendations of the International Scientific Committee: Evaluation and treatment of urinary incontinence, pelvic organ prolapse, and fecal incontinence. *Neurourology* and Urodynamics: Official Journal of the International Continence Society, 29(1), 213-240.
- Nygaard, I., Barber, M. D., Burgio, K. L., Kenton, K., Meikle, S., Schaffer, J., ... & Pelvic Floor

Disorders Network. (2008). Prevalence of symptomatic pelvic floor disorders in US women. *Jama*, *300*(11), 1311-1316.

- 3. Norton, P., & Brubaker, L. (2006). Urinary incontinence in women. Lancet, 367:57-67.
- 4. Blaivas, G. J. (2002). Urinary incontinence: Epidemiology, pathophysiology, evaluation, and management overview. Campbell's Urol, 2:1207-52.
- 5. D. K. (2003). Newman, Stress Urinary Incontinence in Women: Involuntary urine leakage during physical exertion affects countless women. AJN The American Journal of Nursing, 103(8), 46-55.
- Avery, K., Donovan, J., Peters, T. J., Shaw, C., Gotoh, M., & Abrams, P. (2004). ICIQ: a brief and robust measure for evaluating the symptoms and impact of urinary incontinence. *Neurourology and Urodynamics: Official Journal of the International Continence Society*, 23(4), 322-330.
- Dupont, M. C., Albo, M. E., & Raz, S. (1996). Diagnosis of stress urinary incontinence. *Urologic Clinics*, 23(3), 407-416.
- Teleman, P. M., Lidfeldt, J., Nerbrand, C., Samsioe, G., Mattiasson, A., & WHILA Study Group. (2004). Overactive bladder: prevalence, risk factors and relation to stress incontinence in middle-aged women. *BJOG: An International Journal of Obstetrics & Gynaecology*, 111(6), 600-604.
- Dietz, H. P., Hansell, N. K., Grace, M. E., Eldridge, A. M., Clarke, B., & Martin, N. G. (2005). Bladder neck mobility is a heritable trait. *BJOG: An International Journal of Obstetrics* & *Gynaecology*, 112(3), 334-339.
- Frazer, M. I., Haylen, B. T., & Sutherst, J. R. (1989). The Severity of Urinary Incontinence in Women Comparison of Subjective and Objective Tests. *British journal of urology*, 63(1), 14-15.
- Fan, H. L., Chan, S. S. C., Law, T. S. M., Cheung, R. Y. K., & Chung, T. K. H. (2013). Pelvic floor muscle training improves quality of life of women with urinary incontinence: a prospective study. Australian and New Zealand Journal of Obstetrics and Gynaecology, 53(3), 298-304.
- Kashanian, M., Ali, S. S., Nazemi, M., & 12. Bahasadri, S. (2011). Evaluation of the effect of pelvic floor muscle training (PFMT or Kegel exercise) and assisted pelvic floor muscle training (APFMT) by a resistance device (Kegelmaster device) on the urinary incontinence in women "comparison between them: a randomized trial". European Journal of **Obstetrics** x Gynecology and Reproductive Biology, 159(1), 218-223.
- 13. Herzog, A. R., & Fultz, N. H. (1990). Prevalence and incidence of urinary incontinence in community-dwelling populations. *Journal of the American Geriatrics Society*, *38*(3), 273-281.

- Brown, J. S., Grady, D., Ouslander, J. G., Herzog, A. R., Varner, R. E., & Posner, S. F. (1999). Prevalence of urinary incontinence and associated risk factors in postmenopausal women. *Obstetrics* & *Gynecology*, 94(1), 66-70.
- Black, N. A., Bowling, A., Griffiths, J. M., Pope, C., & Abel, P. D. (1998). Impact of surgery for stress incontinence on the social lives of women. BJOG: An International Journal of Obstetrics & Gynaecology, 105(6), 605-612.
- 16. Fultz, N. H., & Herzog, A. R. (1996). Epidemiology of urinary symptoms in the geriatric population. *Urologic Clinics*, 23(1), 1-10.
- Swift, S. E., & Ostergard, D. R. (1995). Evaluation of current urodynamic testing methods in the diagnosis of genuine stress incontinence. *Obstetrics* & *Gynecology*, 86(1), 85-91.