

Uterine Artery Doppler Ultrasound Measurement of Preterm Labor in Pregnant Women with Threatened Preterm Labor

Mahnaz Syed^{1*}, Mohammed Ziaur Rahman², Syed Tasnuv Sami³

¹Junior Consultant (Obs and Gynae), Upazila Health Complex, Golapgonj, Sylhet, Bangladesh

²Senior Consultant (Medicine), 250 Bed General Hospital, Moulvibazer, Bangladesh

³Outdoor Medical Officer (Medicine), Sylhet Women's Medical College Hospital, Sylhet, Bangladesh

DOI: [10.36348/sijog.2023.v06i03.004](https://doi.org/10.36348/sijog.2023.v06i03.004)

Received: 03.02.2023 | Accepted: 08.03.2023 | Published: 11.03.2023

*Corresponding author: Dr. Mahnaz Syed

Junior Consultant (Obs and Gynae), Upazila Health Complex, Golapgonj, Sylhet, Bangladesh

Abstract

Introduction: One of the most common causes of hospitalisation during pregnancy is preterm labour. Nearly one-fifth of women hospitalised for preterm labour. The frequency of preterm births is about 12%–13% in the USA and 5%–9% in many other developed countries. One of the main causes of preterm delivery is preterm premature rupture of membranes, while pre-eclampsia and foetal growth restriction (FGR) can be identified as other common causes that could lead to such complications. **Objective:** to assess the value of uterine artery doppler ultrasound measured of preterm labor in pregnant women with threatened preterm labor. **Methods:** We performed a single-center cohort study was carried out at Obstetrics and Gynecology, Sylhet MAG Osmani Medical College Hospital & Jalalabad Clinic, Modhushohid, Sylhet, Bangladesh from January to December 2020. Total 141 women were selected for the present study after applying inclusion and exclusion criteria. All participating women signed informed written consent. The study included singleton pregnant women who present to the casualty at gestations between 28 and 34 weeks of gestation with symptoms and signs of threatened preterm labor (defined as presence of at least one uterine contraction per 10 minutes, lasting at least 30 seconds, with a cervical dilatation ≤ 3 cm, and a cervical effacement $< 80\%$). On admission, during obstetric ultrasound scanning, bilateral uterine artery Doppler ultrasound velocimetry was performed using the transabdominal technique. Uterine artery Doppler scans were both conducted at the peak of uterine contraction and in between contractions when the uterus is fully relaxed. **Results:** Total 141 women presenting with threatened preterm labor were included in the study. The mean gestational age at presentation was 30.86 ± 1.71 weeks (range: 28 – 33.86 weeks). Among 45 (31.9%) delivered within 7 days, while 96 (68.1%) delivered after 7 days of presentation. The mean uterine artery pulsatility index (UA-PI) measured both basally and at the peak contraction were significantly higher among women who delivered within 7 days. ROC curves showed that both basal and contraction UA-PI were significant predictors of delivery within 7 days. There was a significant negative correlation between contraction UA-PI and birth weight. **Conclusion:** Uterine artery Doppler ultrasound velocimetry measured in women with threatened preterm labor, seems to be a significant predictor of actual preterm labor within 7 days of admission.

Keywords: Preterm Labor, Uterine Artery Doppler, Threatened Preterm Labor.

Copyright © 2023 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

One of the most common causes of hospitalisation during pregnancy is preterm labour [1]. Nearly one-fifth of women hospitalised for preterm labour [2]. The frequency of preterm births is about 12%–13% in the USA and 5%–9% in many other developed countries [3]. One of the main causes of preterm delivery is preterm premature rupture of

membranes, while pre-eclampsia and foetal growth restriction (FGR) can be identified as other common causes that could lead to such complications [3,4]. Several studies have reported an effect of smoking on length of gestation [5, 6]. It is often that following the assessment for preterm labour, the patient is admitted to centres that offer therapies such as tocolysis and steroids to inhibit uterus contractions or alternatively referred to hospitals for further treatments

[7-9]. Preterm birth (PTB), defined as delivery before 37 weeks of gestation, remains a major burden in obstetric care, affecting over 15 million babies worldwide each year [10, 11]. Despite various preventive measures, 1 million neonatal deaths are attributable to complications of PTB, which makes it the leading cause of death in children under the age of 5 years [12]. To reduce these numbers, it is crucial to understand the mechanisms underlying prematurity and develop targeted interventions for prevention. During the normal development of the placental structure, trophoblast cells invade the myometrium and cause remodeling of maternal spiral arteries. These spiral arteries convert to a low-resistance, high-flow state [13], management of such women who present with threatened preterm labor includes exclusion of serious obstetric and non-obstetric conditions that might have precipitated preterm labor, tocolysis, maternal steroid administration to enhance fetal lung maturation and transfer to a tertiary maternity unit with adequate neonatal facilities, should preterm delivery ensue [14]. Deliveries at such a tertiary unit as well as, proper and timely handling of the preterm neonate are both essential for minimizing the risk of neonatal morbidity and mortality [15]. Therefore, prediction of preterm labor among women who present with symptoms of threatened preterm labor is vital. True uterine contractions that can eventually be effective to cause cervical dilatation is associated with compression of the intramyometrial arteries resulting in reduction or even disappearance of the diastolic component of the uterine artery blood flow [16, 17]. Therefore, measurement of uterine artery (UTA) Doppler during contractions may be considered as a predictor of preterm birth in women who are admitted for preterm labour [18, 19].

MATERIALS AND METHODS

We performed a single-center cohort study was carried out at Obstetrics and Gynecology, Sylhet MAG Osmani Medical College Hospital & Jalalabad Clinic, Modhushohid, Sylhet, Bangladesh from January to December 2020. Total 141 women were selected for the present study after applying inclusion and exclusion criteria. All participating women signed informed written consent. The study included singleton pregnant women who present to the casualty at gestations between 28 and 34 weeks of gestation with symptoms and signs of threatened preterm labor (defined as presence of at least one uterine contraction per 10 minutes, lasting at least 30 seconds, with a cervical dilatation ≤ 3 cm, and a cervical effacement $< 80\%$). Women with ruptured fetal membranes, signs of chorioamnionitis, fetal compromise, maternal obstetric or non-obstetric complication (e.g. severe preeclampsia/eclampsia, acute abdomen, liver impairment) were not recruited in the study. After revising medical history and performing general, abdominal and local examination, obstetric ultrasound, non-stress test (NST) for fetal heart rate tracing and complete blood count (CBC), women were subjected to

intramuscular steroid injection (dexamethasone 6 mg every 12 hours for 48 hours) and tocolysis (using oral nifedipine 10 mg capsule every 20 minutes until uterine contractions have faded away). On admission, during obstetric ultrasound scanning, bilateral uterine artery Doppler ultrasound velocimetry was performed using the transabdominal technique.

All ultrasound scans and uterine artery Doppler ultrasound measurements were taken by a sonographer with at least 3-year experience with such scans. After receiving the standard management, women were observed for few days in the hospital. According to the practice guidelines adopted by our Department, no long-term maintenance tocolytic treatment was given to women who recovered from threatened preterm labor. Women who had not delivered imminently were contacted later by phone to note the gestational age at delivery and birth weight.

Statistical Analysis

Data were analysed using SPSS software version 20 (SPSS Inc., Chicago, IL, USA). Independent-samples t-test was used to compare the results for variables. The indices used for this study were the mean, standard deviation and correlation coefficient. Normally distributed variables were described in terms of mean and standard deviation; the difference between two independent groups for such variables was analyzed using independent t-test and expressed in terms of mean difference and its 95% confidence interval. Non-normally distributed variables were described in terms of median and interquartile; the difference between two independent groups for such variables was analyzed using Mann Whitney's U-test and expressed in terms of median difference and its 95% confidence interval. Following the analysis of logistic regression, a receiver operating characteristic curve was used to determine the optimal cut-off values that considered $P < 0.05$ as significant.

RESULTS

Total 141 women presenting with threatened preterm labor were included in the study. The mean age of included women was 32.19 ± 5.14 years (range: 24-41 years). The median parity was 2 (range: 0-4; interquartile range: 1-3). The mean body mass index (BMI) was 27.37 ± 4.7 kg/m² (range: 17.58-39.19 kg/m²). The mean gestational age at presentation was 30.86 ± 1.71 weeks (range: 28-33.86 weeks). Of the included 141, 45 (31.9%) delivered within 7 days, while 96 (68.1%) delivered after 7 days of presentation. There were no significant differences between women of both groups regarding the age, parity, BMI or gestational age (table-1). The mean uterine artery pulsatility index (UA-PI) measured both basally and at the peak contraction were significantly higher among women who delivered within 7 days (table-2). ROC curves showed that both basal and contraction UA-PI were significant predictors of delivery within 7 days (figure-

1). There was a significant negative correlation between contraction UA-PI and birth weight ($r=-0.478$, 95% CI

(-0.563 to -0.383), $p<0.001$) (figure-2).

Table 1: Difference between Groups regarding Initial Characteristics (N=141)

	Women Delivered ≤ 7 Days (n=45)	Women Delivered >7 Days (n=96)	MD/MedD (95% CI)	p-value
Age (years)	32.8 \pm 5.3	31.9 \pm 5.1	1.0 (-3.0 to 5.1)	0.614 1
Parity	2 (1 – 3)	2 (1 – 3)	-1.0 (-2.0 to 0.0)	0.243 2
BMI (kg/m ²)	27.7 \pm 4.5	27.2 \pm 4.8	-0.12 (-3.8 to 3.6)	0.947 1
Gestational Age (weeks)	30.6 \pm 1.7	30.9 \pm 1.7	-0.28 (-1.40 to 0.84)	0.618 1

Data presented as mean \pm standard deviation; or median (interquartile range) BMI body mass index MD (95% CI) mean difference and its 95% confidence interval MedD (95% CI) median difference and its 95%

confidence interval 1 Analysis using independent student’s t-test 2 Analysis using Mann-Whitney’s U-Test.

Table 2: Difference between Groups regarding UA-PI (N=141)

	Women Delivered ≤ 7 Days (n=45)	Women Delivered >7 Days (n=96)	MedD (95% CI)	p-value
Basal UA-PI	0.83 (0.76 – 0.9)	0.78 (0.71 – 0.84)	0.05 (0.03 to 0.07)	<0.001
Contraction UA-PI	1.91 (1.58 – 2.17)	1.11 (0.91 – 1.35)	0.74 (0.64 to 0.85)	<0.001

Data presented as median (interquartile range) UA-PI uterine artery pulsatility index MedD (95% CI)

median difference and its 95% confidence interval 1 Analysis using Mann-Whitney’s U-Test.




Fig. 1: ROC Curves for Basal and Contraction UA-PI in Prediction of Preterm Labor within 7 Days.

Table 3: Validity of cut-offs.

	AUC	Cutoff Value	Specificity	Specificity	LR+	LR
Basal UA-PI	0.651 (0.592 to 0.707)	64.84 (54.1 to 74.6)	55.73 (48.4 to 62.9)	55.73 (48.4 to 62.9)	1.46 (1.2 to 1.8)	0.63 (0.5 to 0.9)
Contraction UA-PI	0.892 (0.850 to 0.925)	76.92 (66.9 to 85.1)	91.67 (86.8 to 95.2)	91.67 (86.8 to 95.2)	9.23 (5.7 to 15.0)	0.25 (0.2 to 0.4)

Data presented as value (95% confidence interval) UA-PI uterine artery pulsatility index AUC area under the curve LR+ positive likelihood ratio LR-

negative likelihood ratio $r = -0.478$, 95% CI (-0.563 to -0.383), $p < 0.001$.




Fig. 2: Scatterplot for the Association between Contraction UA-PI and Birth Weight.

DISCUSSION

Our study has both strengths and limitations. A major strength of this study is that we included a cohort with a large sample size. We recruited women with both a low-risk (midwifery practices) and high risk (tertiary center) for spontaneous PTB. Data were derived from a local ultrasound registry (ASTRAIA), in which both data on delivery and other pregnancy outcomes are registered. All ultrasound scans were conducted by certified sonographers. Although the current evidence assigns the cervical length as an objective method for prediction of preterm labor [5, 20], transvaginal scan seems undesirable to some pregnant women. In addition, each of UA Doppler scan and cervical length measurement assesses a different component of preterm labor; the former assesses the contraction component, while the latter assesses cervical ripening [21], and therefore, each of them can identify women at risk of preterm labor, yet not showing changes in the other component. Uterine artery Doppler scan is, however, disadvantaged by the fact that its indices should be affected in women with abnormal trophoblast invasion or placental insufficiency, and in women with multiple pregnancy or polyhydramnios due to rise in intrauterine pressure [12]. The current study showed a promising role of uterine artery Doppler measurement in prediction of actual preterm labor in women presenting with preterm labor. A total 141 women presenting with threatened preterm labor were included in the study. The mean age of included women was 32.19 ± 5.14 years (range: 24-41 years), found no significant difference between both groups regarding the screening uterine artery Doppler

findings performed at 18–24 weeks of gestation [22], these findings does not contradict the results of the current study for two obvious reasons. First, that retrospective analysis targeted asymptomatic pregnant women who had no threatened preterm labor. Second, the timing and circumstances of uterine artery Doppler scanning are both quite different in what proposed in the current study. The main value of uterine artery Doppler is in women with threatened preterm labor. This is further confirmed by two studies conducted on the role of uterine artery Doppler velocimetry in prediction of spontaneous preterm labor in women with threatened preterm labor. In one of them that was conducted by Olgan and Celiloglu (2016), the authors studied the relationship between contraction-based uterine artery PI and preterm delivery in 172 women with threatened preterm labor at gestations between 24 and 32 weeks [12]. The mean values of basal and contraction PI among women who delivered within 7 days were significantly higher than those in women who delivered after 7 days [0.76 ± 0.22 vs. 0.98 ± 0.37 , $p < 0.001$; and 2.66 ± 1.05 vs. 1.03 ± 0.64 , $p < 0.001$; respectively and respectively]. In this study, cervical length was also evaluated. The authors found that accuracy of uterine artery PI was comparable to that of cervical length measured by transvaginal scan in prediction of preterm labor within 7 days [AUC 0.88, 95% CI (0.82 to 0.94), $p < 0.05$ and 0.85, 95% CI (0.77 to 0.93), $p < 0.05$, respectively] [12]. In the second very recent study, just published in November 2019, conducted on 100 Iranian pregnant women admitted for having threatened preterm labor at a tertiary maternity hospital in Tehran, the uterine artery PI was

significantly higher among women who delivered within 48 hours [than those who delivered after 48 hours: 1.4 ± 0.5 vs. 0.9 ± 0.3 , respectively, $p<0.001$], women who delivered within 7 days [than those who delivered after 7 days: 1.1 ± 0.3 vs. 0.9 ± 0.3 , respectively, $p=0.01$], and women who delivered within 14 days [than those who delivered after 14 days: 1.3 ± 0.3 vs. 0.9 ± 0.3 , respectively, $p=0.001$]. In this study the best cutoff values for uterine artery PI for prediction of labor within 48 hours, 7 days and 14 days were 0.98 [sensitivity 75% and specificity 70%], 1.0 [sensitivity 84% and specificity 76%] and 0.95 [sensitivity 63% and specificity 69%], respectively [23]. The figures reported by this study are quite different from those of the current study and those reported by Olgan and Celiloglu [21]. This might be explained by relatively small sample size and the measurement of uterine artery Doppler in the relaxed state of the uterus.

CONCLUSION

In conclusion, uterine artery Doppler ultrasound velocimetry measured in women with threatened preterm labor, seems to be a significant predictor of actual preterm labor within 7 days of admission.

Conflict of Interest: None.

Source of Fund: Nil.

REFERENCES

- Rundell, K., & Panchal, B. (2017). Preterm labor: Prevention and management. *Am Fam Physician*, 95, 366-72.
- Sentilhes, L., Senat, M. V., Ancel, P. Y., Azria, E., Benoist, G., Blanc, J., ... & Langer, B. (2016). Prevention of spontaneous preterm birth (excluding preterm premature rupture of membranes): Guidelines for clinical practice-Text of the Guidelines (short text). *Journal de gynécologie, obstétrique et biologie de la Reproduction*, 45(10), 1446-1456.
- Goldenberg, R. L., Culhane, J. F., Iams, J. D., & Romero, R. (2008). Epidemiology and causes of preterm birth. *Lancet*, 371, 75-84.
- Keelan, J. A., & Newnham, J. P. (2017). Recent advances in the prevention of preterm birth. *F1000Res*, 6, pii: F1000 Faculty Rev-1139.
- Ion, R., & Bernal, A. L. (2015). Smoking and preterm birth. *Reprod Sci.*, 22, 918-26.
- Chan, A., Keane, R. J., & Robinson, J. S. (2001). The contribution of maternal smoking to preterm birth, small for gestational age and low birthweight among aboriginal and non-aboriginal births in South Australia. *Med J Aust*, 174, 389-93.
- Hollingworth, J., Pietsch, R., Epee-Bekima, M., & Nathan, E. (2018). Time to delivery: Transfers for threatened preterm labour and prelabour rupture of membranes in Western Australia. *Aust J Rural Health*, 26, 42-7.
- Menon, R., & Richardson, L. S. (2017). Preterm prelabor rupture of the membranes: A disease of the fetal membranes. *Semin Perinatol*, 41, 409-19.
- Park, C. W., Park, J. S., Moon, K. C., Jun, J. K., & Yoon, B. H. (2016). Preterm labor and preterm premature rupture of membranes have a different pattern in the involved compartments of acute histologic chorioamnionitis and/or funisitis: Patho-physiologic implication related to different clinical manifestations. *Pathol Int*, 66, 325-32.
- Gazmararian, J. A., Petersen, R., Jamieson, D. J., Schild, L., Adams, M. M., Deshpande, A. D., & Franks, A. L. (2002). Hospitalizations during pregnancy among managed care enrollees. *Obstetrics & Gynecology*, 100(1), 94-100.
- Fonseca, E., Yu, C. K. H., Singh, M., Papageorgiou, A. T., & Nicolaides, K. H. (2006). Relationship between second-trimester uterine artery Doppler and spontaneous early preterm delivery. *Ultrasound in obstetrics & gynecology*, 27(3), 301-305.
- Fleischer, A., Anyaegbunam, A. A., Schulman, H., Farmakides, G., & Randolph, G. (1987). Uterine and umbilical artery velocimetry during normal labor. *American journal of obstetrics and gynecology*, 157(1), 40-43.
- JANBU, T., & NESHETM, B. T. (1987). Uterine artery blood velocities during contractions in pregnancy and labour related to intrauterine pressure. *BJOG: An International Journal of Obstetrics & Gynaecology*, 94(12), 1150-1155.
- Ingemarsson, I., & Lamont, R. F. (2003). An update on the controversies of tocolytic therapy for the prevention of preterm birth. *Acta obstetrica et gynecologica Scandinavica*, 82(1), 1-9.
- Sotiriadis, A., Papatheodorou, S., Kavvadias, A., & Makrydimas, G. (2010). Transvaginal cervical length measurement for prediction of preterm birth in women with threatened preterm labor: a meta-analysis. *Ultrasound Obstet Gynecol.*, 35, 54-64.
- Strigini, F. A., Lencioni, G., De Luca, G., Lombardo, M., Bianchi, F., & Genazzani, A. R. (1995). Uterine artery velocimetry and spontaneous preterm delivery. *Obstetrics & Gynecology*, 85(3), 374-377.
- Oskovi Kaplan, Z. A., & Ozgu-Erdinc, A. S. (2018). Prediction of Preterm Birth: Maternal Characteristics, Ultrasound Markers, and Biomarkers: An Updated Overview. *Journal of pregnancy*. Article ID 8367571, 8 pages. <https://doi.org/10.1155/2018/8367571>
- Parra-Cordero, M., Sepúlveda-Martínez, A., Rencoret, G., Valdés, E., Pedraza, D., & Muñoz, H. (2014). Is there a role for cervical assessment and uterine artery Doppler in the first trimester of pregnancy as a screening test for spontaneous preterm delivery?. *Ultrasound in Obstetrics & Gynecology*, 43(3), 291-296. Cobian-Sanchez, F., Prefumo, F., Bhide, A., & Thilaganathan, B.

- (2004). Second-trimester uterine artery Doppler and spontaneous preterm delivery. *Ultrasound in Obstetrics and Gynecology: The Official Journal of the International Society of Ultrasound in Obstetrics and Gynecology*, 24(4), 435-439.
19. Mella, M. T., & Berghella, V. (2009). Prediction of preterm birth: cervical sonography. *Semin Perinatol*, 33, 317-324.
20. Cobian- Sanchez, F., Prefumo, F., Bhide, A., & Thilaganathan, B. (2004). Second- trimester uterine artery Doppler and spontaneous preterm delivery. *Ultrasound Obstet Gynecol.*, 24, 435- 9.
21. Olgan, S., & Celiloglu, M. (2016). Contraction-based uterine artery Doppler velocimetry: novel approach for prediction of preterm birth in women with threatened preterm labor. *Ultrasound Obstet Gynecol.*, 48, 757-764.
22. Scott, C. L., Chavez, G. F., Atrash, H. K., Taylor, D. J., Shah, R. S., & Rowley, D. (1997). Hospitalizations for severe complications of pregnancy, 1987–1992. *Obstetrics & Gynecology*, 90(2), 225-229.
23. Naemi, M., Eslamian, L., Teimoory, N., Moshfeghi, M., & Tajik, A. (2019). Predicting preterm labour by means of uterine artery doppler velocimetry during peak uterine contraction in patients with normal cervical length. *Advances in Human Biology*, 9(1), 61-64.