

Clinical profiles and Outcome of External DCR and TC-LASER DCR

Syeed Mehub Ul Kadir^{1*}, Azizur Rahman Alam², Siddiqur Rahman³, Abid Akbar⁴, Ummay Kawsar⁵, Md. Tauhidur Rahman⁶, Jayanta Kumar Das⁷

¹Assistant Professor, Sheikh Fajilatunnesa Mujib Eye Hospital and Training Institute, Bangladesh

²Consultant, Bangladesh Eye Hospital and Institute, Dhaka, Bangladesh

^{3,4}Consultant, Vision Eye Hospital, Dhaka, Bangladesh

⁵Associate Professor, MH Samarita Medical College, Dhaka Bangladesh

⁶Associate Professor, Shahid Ziaur Rahman Medical College, Bogra Dhaka Bangladesh

⁷Senior Consultant, The Eye Care Center, Guwahati, Assam, India

DOI: [10.36348/sjbr.2021.v06i06.001](https://doi.org/10.36348/sjbr.2021.v06i06.001)

| Received: 13.04.2021 | Accepted: 22.05.2021 | Published: 02.06.2021

*Corresponding author: Syeed Mehub Ul Kadir

Abstract

Aim: To assess the anatomical and functional outcome of transcanalicular LASER DCR compared to external DCR.

Methods: A quasi study had been carried out in two tertiary eye hospitals of Bangladesh from January 2016 to December 2020. Group A included all patients selected for external DCR, and group B had been selected for transcanalicular laser DCR. Variable included age, gender, anatomical outcome, functional outcome, and surgery-related complications. Statistical analysis had been done by Quick Calcs Graph Pad software. **Results:** The total evaluated patients were 119 patients in group A and 46 patients in group B. The anatomical success rate was 93.3% in group A and 84.8% in group B. The functional success rate was 86.5% in group A and noted 84.8% in group B. Minimal skin scar was observed after six weeks of surgery in 98 (82.3%) cases of group A. **Conclusion:** The anatomical success rate is higher in external DCR, but the functional outcomes are almost the same in both groups.

Keywords: Dacryocystitis, nasolacrimal duct obstruction, external DCR, Transcanalicular LASER DCR, Anatomical and functional outcome.

Abbreviations: DCR= Dacryocystorhinostomy, TC= Transcanalicular, NLD=Nasolacrimal duct, PNADO= Primary acquired nasolacrimal duct obstruction, MMC= Mitomycin-C.

Copyright © 2021 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

A Dacryocystorhinostomy (DCR) surgery is making an anastomosis between the lacrimal sac and the nasal cavity at the level of the middle meatus by cutting the intervening bone. This new opening is proximal to the site of nasolacrimal duct obstruction and re-establishes the tear flow into the nose. Different approaches are available for DCR surgery, e.g. external, transnasal and both. These approaches include external or conventional DCR, Non LASER endoscopic DCR, endoscopic endonasal laser DCR, and transcanalicular laser-assisted DCR. The traditional or external DCR is considered as the standard gold technique for managing acquired nasolacrimal duct obstruction [1, 2]. Caldwell first introduced the transnasal DCR in 1893 but did not widely accept it due to complex visualization of the nasal cavity and perioperative bleeding [3]. With the advancement of endoscopic equipment, the endoscopic endonasal approach had popularized with a reasonably good outcome. The LASER assisted endoscopic

approach had revolutionized DCR surgery, especially for cosmetic concern, precise ostium, haemostasis, and less surgical morbidity [1-2, 4-7]. Different types of LASER are used in DCR surgery and most useful with minor collateral damage. Diode laser-assisted DCR included both endoscopic and external approaches and offers many advantages over other LASER DCR and conventional DCR [4-6,8]. Skin incision sparing DCR is the current mainstay of managing congenital and acquired nasolacrimal duct obstruction for young children and adults. We assessed the surgical strategies and compared the outcome of LASER DCR and conventional DCR.

Patients and Methods: This quasi interventional study had been carried out in Bangladesh eye hospital and institute of Dhaka, Bangladesh, and Vision eye hospital, Dhaka, Bangladesh. We started the research in January 2016 and completed it on 30 June 2020. Pre-operative ophthalmic and nasal cavity

evaluation and pre-anaesthetic check-up had made in all cases. All cases were divided into two groups; group A and Group B. In Group A, all patients underwent external or conventional dacryocystorhinostomy (External DCR). Group B included all patients who had managed transcanalicular LASER dacryocystorhinostomy (TC-DCR). External DCR was used for all the patients with Failed DCR. External DCR and Transcanalicular LASER DCR had offered with counselled potential advantages and disadvantages of surgical procedures for all the cases of primary acquired nasolacrimal duct obstruction. TC LASER DCR was costly than external DCR. In our study, the lowest age was 12 years, and the highest was 86 years. This study excluded all the patients suspected of lacrimal neoplasm, rhinosporidiosis of the lacrimal sac, nasal neoplasm. Anatomical success had assessed by the patency of the lacrimal passage on irrigation with normal saline. The operational success had been evaluated by the absence or insignificant epiphora without any ocular and eyelid diseases. Data were collected and analyzed by Graph Pad Quick Calcs Software.

SURGICAL TECHNIQUES

Anaesthesia: Most of the patients were operated by local anaesthesia (LA) with intravenous sedation; only two cases of group A were operated by general anaesthesia. We had used a mixture of Hyaluronidase (1500IU) mixed with bupivacaine HCL 0.5% (5 mg/ml) and lidocaine (2%) with epinephrine (0.0005%) as LA. We used plain lidocaine (2%) for hypertensive patients with chronic dacryocystitis. The LA had been injected as Infratrochlear nerve block, infraorbital nerve block, ethmoidal nerve block and dorsal nasal nerve block for DCR. Intravenous sedation with 1 to 2 ml of Midazolam 1 mg/ml and Fentanyl 0.5 to 2 mcg/kg over 1-2 minutes. We sprayed 10% lignocaine solution in the nasal cavity to reduce the sensitization of nasal mucosa. In all cases, a qualified anaesthetist was present during surgery to administer intravenous drugs and monitor the patients' vitals.

Nasal Packings: a 10-15cm ribbon Gause socked with 2% Lignocaine jelly, oxymetazoline nasal drop, Inj. Adrenaline 1 ml and introduced as a posterior nasal pack throughout the surgery, and also introduced an anterior nasal packing (3-4 cm) to the middle meatus

at least 5 minutes to taught nasal mucosa and also for hemostasis purpose as nasal packings.

TC LASER DCR: The TC LASER DCR system includes a 980 nm wavelength Diode LASER with a 600 μ m fibre optic probe, 0° angle rigid camera-mounted nasal endoscope. The LASER fibre optic probe was used for this procedure through canaliculi to the sac. After punctual dilatation with Nettleship punctum dilator, the laser probe was inserted horizontally into the sac through the upper punctum and canalicular system and then advanced obliquely (about 60° to 70°) vertically downward, medially and backwards, nearly the same as in lacrimal probing. Then, the probe had been pushed till felt a stiff resistance was along the nasolacrimal duct to the lateral wall of the nasal cavity. A 4 mm diameter, 20 cm long 0° angled rigid camera-mounted nasal endoscope was introduced into the nasal cavity to visualize the Laser glow of the pilot beam. The properly focused red light glow of laser (pilot) beam in the middle meatus (Figure 1a). The LASER glow will reveal the thinnest portion of the lacrimal bone, which is anterior and inferior to the insertion of the middle turbinate. The middle turbinate medialization is vital for good exposure and protection from LASER heat. A continuous contact mode of diode laser with 980 nm wavelength had been used to create a nasolacrimal osteotomy by ablating the bone and mucosal tissues by pushing the beam towards the nasal cavity applying 3-4 watt of power. Both the pilot beam and 980 nm delivered Laser energy through the same LASER optical fibre. This procedure was repeated through the lower punctum and canaliculi to extend the ostium. The osteotomy was enlarged up to 7-8 mm vertically and 5 mm horizontally by pulling up followed by pushing down the laser probe in a see-saw movement (Figure 1b). A bi-canalicular silicone lacrimal stent was introduced through both canaliculi (Figure -2) and fixed to the medial wall of anterior nares in all cases, and kept in situ up to 6 weeks of surgery. After removing all nasal packing, a piece of merocel pack (compressed dehydrated sponge composed of hydroxylated polyvinyl acetate) was introduced into the space between the middle turbinate and newly created osteotomy to prevent adhesion of middle turbinate and also to prevent the postoperative hemostasis and kept it for seven days.




Fig-1a-B: The LASER glow is showing through the thinnest portion of the lacrimal bone, b. An osteotomy is created at the level of middle meatus by a multimode diode laser beam. **Figure 2:** Intubation of bicanalicular silicone DCR tube after LASER DCR

External DCR: A J-shaped incision was given to all cases to achieve minimal or no skin scar postoperatively. Dissection had made and identified the medial palpebral ligament, making a lacrimal mucosal flap, then created a bone osteotomy by cutting the intervening bone. The nasal mucosal flap had prepared and made an anastomosis between the nasal and lacrimal mucosal flap by 6-0 vicryl (Figure-3). Used Mitomycin C (0.02%), particularly in between the

mucosal and lacrimal flaps with a surgical sponge/cotton pledge for 3 minutes and then rinsed. MMC had been used in patients who had excessive granulation tissue at the surgical site. Silicone intubation was introduced in all cases and kept in the nasal cavity for six weeks of surgery. We placed a nasal pack with antibiotic ointment at the end of the surgery for 24 hours.



Fig-3: Steps of External DCR: J-Shaped incision with desection of underlying tissue with radi frequency cautery, Exposure of medial palpebral ligament, expose the fossa for the lacrimal sac after making a mucosal flap, creating a nasal mucosal flap, intubation of a DCR tube, an anastomosis of Lacrimal sac mucosa and Nasal mucosal flap in external DCR, and closing the wound with 6-0 vicryl.

RESULTS

A total of 165 patients were evaluated in this study including male (84 cases, 50.9%), and female (81 cases, 49.09%). The over all mean age was 50.9 years. Primary acquired nasolacrimal duct obstruction was found (PANDO) on sac patency test in 121 (73.3%) cases, and others (26.7%) was associated with failed DCR. patients. In group A, the total number of patients was 119, with 56.3% male and 43.7% female. 75 (63%) patients presented with PANDO, and 44 (37%) patients presented with failed DCR. In 119 patients, Comorbidities were in 77 (64.7%) patients. 24 (20.2%) Patients had taken anticoagulant drug like Ecospirin, Clopidogrel. The age range was 12 years to 86 years and the mean age 57.32 years. In Group B, all 46

patients had presented with PANDO. Comorbidities were associated in only 10 (21.7%) cases. The female was 29 (63%) cases, and the male was 17 (37%). The age ranges from 24 years to 69 years, and the mean age was 44.59 years. The mean operating time was 43.52 minutes in group A and 22.19 minutes in group B patients. The anatomical success rate had been found in 111 cases (93.3%) of group A. Though, the functional success rate had been noted in 103 (86.5%) cases of group A in one year follow up time (Table-1). The anatomical and functional success rate was observed in 39 (84.8%) patients who managed by TC-LASER DCR (Group B). A Sign and binomial test had been calculated and the P-value was highly significant (<0.0001) in both groups.

Table-1: Distribution of Demographic profiles, clinical profiles and outcome values of both groups

Variables	Group-A	Group-B
Demographic profiles:		
Age Range (Year)	12 to 86	24 to 69
Mean Age (Year)	57.32	44.59
Male	67 (56.3%)	17 (37%)
Female	52 (43.7%)	29 (63%)
Clinical Profile:		
PANDO	75 (63%)	46 (100%)
Failed DCR	44 (37%)	00 (0%)
Comorbidities	77 (64.7%)	10 (21.7%)
H/o Anticoagulant drug	24 (20.2%)	08 (17.4%)
Outcomes:		
Mean surgery time	43.52 minutes	22.19 minutes
Anatomical Success	111 (93.3%)	39 (84.8%)
Functional Success	103 (86.5%)	39 (84.8%)
Anatomical Failure	08 (6.7%)	07 (15.2%)
Functional Failure	16 (14%)	07 (15.2%)

In group A, anatomical success and functional success were observed in 97.3% and 93.3% patients respectively who had presented with PANDO. The ultimate functional outcome was achieved in 75% cases who underwent for re-DCR (Table-2). Faint or minimal skin scar was noted in 98 (82.3%) cases after six weeks of external DCR surgery (Figure-4) but reduced to only 12% after three months of surgery. No skin scar in the instances of LASER DCR surgery (Figure-5). One wound dehiscence following external DCR had

managed with conservative treatment followed by secondary wound closure. Minimal postoperative nasal bleeding had noted in 21 (17.64%) cases of group A and 2.17% cases of group B. Complained moderate postoperative pain was up to 4 days of surgery in Group A and two days in group A patients. Felt minimal pain up to 10 days of surgery in group A and up to 7 days in group B patients. There was no scarring on the skin, wound dehiscence in group B patients.

Table-2: Distribution of the outcomes among different clinical entities of the group A patients

Ext DCR,	No. (%)	Anatomical success	Functional Success	Anatomical Failure	Functional Failure
PANDO	75	73 (97.3%)	70 (93.3%)	02(2.67%)	05 (6.7%)
Failed DCR	44	38 (86.4%)	33 (75%)	06 (13.6%)	11 (25%)

The anatomical failure rate was 6.7% in external DCR cases (group A) and 15.2% in TC-LASER DCR cases (group B). The functional failure rate of group A (14%) is near to TC-LASER DCR (15.2%). In group A, inefficient lacrimal pump mechanism was one of the cause of fuctional failure, but stenosis of the ostium or closure of the ostium was the main cause of failure. The success rate depends on patients co-operation during surgery, the clinical condition of the lacrimal drainage system and nasal cavity, surgical experiences, instrumental facilities, pre-operative evaluation and management, and

comorbidities. Per-operative bleeding was more in hypertension and ischaemic heart disease (IHD) patients taking Anti-coagulant medication. LASER DCR was usually selected for the cases of primary acquired nasolacrimal duct obstruction, especially in the younger age group and those who were sensitive to a cosmetic concern. However, 10 cases (21.7%) of older adults and comorbidity patients with PANDO had been operated by LASER DCR to drain the tear from the eye to nasal cavity with minimal surgical trauma and minimum operative time.



Fig-4: Minimal skin scar at the incision site after 7 days of external DCR, and 6 weeks after External DCR surgery. Figure 5: No skin scar after LASER DCR surgery, and intubation in situ after 6 weeks of TC LASER DCR

DISCUSSION

External DCR is a highly successful and gold standard operation for nasolacrimal duct obstruction (NLDO). It is also an effective procedure in revision surgeries for all types of failed DCR cases [9-12]. In recent days, minimally invasive techniques and new technology-based endoscopic approaches have reported high success rates [13-17]. Both Endoscopic endonasal DCR and Transcanalicular LASR DCR procedures are the choice of surgery to avoid skin scar. There is no possibility for skin scarring, wound infection or wound dehiscence. These procedures require additional high-cost surgical equipment and visual systems and need experience in endoscope handling. Skin incision sparing LASER DCR or Endoscopic DCR is helping to preserve the lacrimal pump function by keeping the medial canthal tendon and canalicular system intact.

Having minimal perioperative bleeding rates, short duration of surgery times and quick rehabilitation times [18-21]. Transcanalicular LASER DCR is a safe and fast operative procedure with low morbidity and well-tolerated in primary acquired nasolacrimal duct obstruction. Compared to External DCR, Transcanalicular LASER DCR could do under local anaesthesia with intravenous sedation. It involves precise cutting and removal of bone, lacrimal, and nasal mucosa by ablation and creating a new opening. It is almost bloodless, less time-consuming DCR surgery, leaves no skin scars, preserves ligaments and muscles of the internal canthus, and keeps physiological lacrimal pump function. TC-laser DCR causes minimum pain and minimum nasal bleeding [13, 19, 22, 23].

The success rate of external DCR has been reported from over 89% to 98% [10, 11, 24-26]. The reported success rates of transcanalicular LASER DCR vary from 52% to 96% [18, 19, 22, 26-29]. The surgical success rate are 52%, 56%, 64%, 76%, and 88% in the age group of 20-30 years, 31-40 years, 41-50 years, 51-60 years, and 61-70 years respectively among the patients who underwent transcanalicular laser DCR with silicone tube intubations. The overall success rate is 67% [31]. The mean age was 44.59 years of transcanalicular LASER DCR (group B) in our study. The functional success rate of transcanalicular LASER DCR has been reported from 68% to 80% [8, 32-35]. Recent studies have reported that the success rate in transcanalicular laser-assisted DCR with intubations ranges from 73.3% to 94.2% [36]. There are many causes for the failure of LASER DCR. Common causes are stenosis and scarring of the ostium, fibrosis at the new ostium, membrane formation over the new ostium, and canalicular stenosis resulting in obstruction of the nasolacrimal pathway [9, 10]. The anatomical success is 97.3% of external DCR among the patients of primary NLD obstruction and 84.8% in transcanalicular LASER DCR in this study. The functional success rate is 93.3% of external DCR and 84.8% of LASER DCR in the cases of PANDO. The overall anatomical and functional success rate of external DCR is 93.3% and 86.5%, respectively. The operational success rate was higher in primary external DCR (93.3%) than external re-DCR (75%). The overall anatomical success rate was 85% in external re-DCR [37], but our success rate is 86%. There is no significant difference statistically between the functional success rate of external DCR and transcanalicular LASER DCR [34]. Failure of transcanalicular LASER DCR is occurred due to smaller osteotomy compared to external DCR and the fibrovascular proliferation, which may cause stenosis and scaring off new ostium, especially in the younger age group [31]. New techniques and modifications have been made, such as mitomycin-C intraoperatively in LASER-DCR to reduce the formation of fibrovascular proliferation, which increases the success rate up to 93% [22]. Because the number of fibroblasts decreases or the fibroblasts degenerate with age, which results in less scar tissue formation, the adhesions between the middle turbinate and new osteotomy are among the causes of the failure of LASER DCR [28,38-42]. Strong expression of nasal mucosal heat shock protein 47 also leads to the formation of fibrosis and scar tissue in the young adult patient, which decreases the success rate of LASER DCR [41]. We used a merocel nasal pack between the middle turbinate and the new osteotomy site to prevent adhesion and haemostasis. The osteotomy size was 11.84 mm in diameter at external DCR surgery, but it is reduced to the average size of 1.8 mm by ultrasonic assessment after six months of external DCR surgery [43-44]. We performed a transcanalicular LASER DCR approach due to its better surgical outcome, and LASER can apply directly to the obstructed site. We present our experience of

transcanalicular LASER-assisted-DCR using 980 nm diode lasers using fibre-optic cable was used because it offers high absorption in water and oxyhemoglobin, with very efficient vaporization of bone and soft tissue, and achieves almost bloodless DCR surgery. The new osteotome was created just anterior and inferior to the middle turbinate.

Bone fractures heal more quickly in the younger patient than in older patient due to higher osteoblastic activity. Those mentioned above were the possible factors to reduce the satisfactory laser DCR due to the smaller osteotomy size [31]. In our study, the minimum osteotomy size was 10 mm in length and 10 mm wide in external DCR, and maximum of 8 mm in length and 5 mm in width in transcanalicular LASER DCR. The success rate was higher in external DCR due to the larger osteotomy size. With increasing age, diminished microcirculation contributes to poor tissue regeneration in older patients. The mean operative time was 17.41 minutes in transcanalicular LASER DCR and 49.49 minutes in external DCR [26]. This study showed that the mean surgery time 43.52 minutes in group A (external DCR) and 22.19 minutes in group B (LASER DCR) patients. Silicone intubation at least six weeks helps increasing the success rate of both external DCR and Transcanalicular LASER DCR in our cases. A recent study reported no significant difference between the removal of silicone intubation after two weeks and six weeks of DCR surgeries [45]. Current ongoing investigations will further clarify the efficacy of these newer techniques and modification of surgery. Using mitomycin-C, silicone intubation, and a piece of merocel nasal pack postoperatively are likely to increase the success rate of DCR. The advantages of external DCR includes high success rates due to large osteotomy and can be used it for revision surgery after failed DCR. The Success rate is higher in older age rather than younger age due to high fibroblastic activity. We have recently performed the transcanalicular LASER DCR in paediatric NLD obstruction, extreme older age, and revision surgeries after failing DCR.

CONCLUSIONS

Transcanalicular LASER DCR is a viable surgical option with minimal hazards to external DCR and overall good surgical outcome in primary nasolacrimal duct obstruction. External DCR is still the best treatment option for revision surgeries of failed DCR. Few modifications of surgery and advancement of instruments are helping to achieve the greater success rate of LASER DCR.

REFERENCES

1. Massaro, B. M., Gonnering, R. S., & Harris, G. J. (1990). Endonasal laser dacryocystorhinostomy: a new approach to nasolacrimal duct obstruction. *Archives of ophthalmology*, 108(8), 1172-1176.

2. Weber, R. K., Keerl, R., Schaefer, S. D., & Rocca, R. C. (Eds.). (2007). Atlas of lacrimal surgery. Springer Science & Business Media.
3. Caldwell, G. W. (1893). Two new operations for obstruction of the nasal duct, with preservation of the canaliculi. *Am J Ophthalmol*, 10, 189-192.
4. Hehar, S. S., Jones, N. S., Sadiq, S. A., & Downes, R. N. (1997). Endoscopic holmium: YAG laser dacryocystorhinostomy-safe and effective as a day-case procedure. *The Journal of Laryngology & Otolaryngology*, 111(11), 1056-1059.
5. Reifler, D. M. (1993). Results of endoscopic KTP laser-assisted dacryocystorhinostomy. *Ophthalmic plastic and reconstructive surgery*, 9(4), 231-236.
6. Sadiq, S. A., Hugkustone, C. E., Jones, N. S., & Downes, R. N. (1996). Endoscopic holmium: YAG laser dacryocystorhinostomy. *Eye*, 10(1), 43-46.
7. Seppä, H., Grenman, R., & Hartikainen, J. (1994). Endonasal CO₂-Nd: YAG laser dacryocystorhinostomy. *Acta ophthalmologica*, 72(6), 703-706.
8. Uludag, G., Yeniad, B., Ceylan, E., Yildiz-Tas, A., & Kozer-Bilgin, L. (2015). Outcome comparison between transcanalicular and external dacryocystorhinostomy. *International journal of ophthalmology*, 8(2), 353.
9. Taşkıran, Ç.A. (2014). *Lasers Surg Med*, 46(4):275-80.
10. Taret, K. J., & Custer, P. L. (1995). External dacryocystorhinostomy: surgical success, patient satisfaction, and economic cost. *Ophthalmology*, 102(7), 1065-1070.
11. Buttanri, I. B., Serin, D., Karslıoğlu, S., Akbaba, M., Fazıl, K., Acar, B., & Sevim, M. S. (2012). The outcome of silicone intubation and tube removal in external dacryocystorhinostomy patients with distal canalicular obstruction. *European journal of ophthalmology*, 22(6), 878-881.
12. Buttanri, I. B., Ayıntap, E., Serin, D., Akbaba, M., & Karslıoğlu, S. (2014). Comparison of revision surgeries with transcanalicular diode laser and external approaches in cases with failed transcanalicular diode laser dacryocystorhinostomy. *Ophthalmic Plastic & Reconstructive Surgery*, 30(3), 209-211.
13. Lee, S., & Yen, M. T. (2011). Laser-assisted dacryocystorhinostomy: a viable treatment option?. *Current Opinion in Ophthalmology*, 22(5), 413-418.
14. Karim, R., Ghabrial, R., Lynch, T. F., & Tang, B. (2011). A comparison of external and endoscopic endonasal dacryocystorhinostomy for acquired nasolacrimal duct obstruction. *Clinical ophthalmology (Auckland, NZ)*, 5, 979.
15. Emmerich K.H, Ungerechts R, Rosenberg H.W.M. Possibilities and limits of minimal invasive lacrimal surgery. *Orbit*. 2000; 19(2):67-71.
16. Emmerich, K. H., Lüchtenberg, M., Meyer-Rüsenberg, H. W., & Steinhauer, J. (1997). Dacryocystorhinostomy and laser dacryoplasty: technique and results. *Klinische Monatsblätter für Augenheilkunde*, 211(6), 375-379.
17. Javed Ali, M., & Honavar, S. G. (2012). Assessment of patient satisfaction following external versus transcanalicular dacryocystorhinostomy. *Current eye research*, 37(9), 853-853.
18. Drnovšek-Olup, B., & Beltram, M. (2010). Transcanalicular diode laser-assisted dacryocystorhinostomy. *Indian journal of ophthalmology*, 58(3), 213.
19. Gras-Cabrerizo, J. R., Montserrat-Gili, J. R., León-Vintró, X., Lopez-Vilas, M., Rodríguez-Álvarez, F., Bonafonte-Royo, S., & Masegur-Solench, H. (2013). Endonasal endoscopic scalpel-forceps dacryocystorhinostomy vs endocanalicular diode laser dacryocystorhinostomy. *European journal of ophthalmology*, 23(1), 7-12.
20. Sharma, V., Martin, P. A., Bengler, R., Kourt, G., Danks, J. J., Deckel, Y., & Hall, G. (2005). Evaluation of the cosmetic significance of external dacryocystorhinostomy scars. *American journal of ophthalmology*, 140(3), 359-e1.
21. Ng, D. S. C., & Chan, E. (2016). Techniques to minimize skin incision scar for external dacryocystorhinostomy. *Orbit*, 35(1), 42-45.
22. Hong, J. E., Hatton, M. P., Leib, M. L., & Fay, A. M. (2005). Endocanalicular laser dacryocystorhinostomy: analysis of 118 consecutive surgeries. *Ophthalmology*, 112(9), 1629-1633.
23. Holak S.A, Wysocki M, Holak N, Holak H. Endocanalicular laser dacryocystorhinostomy (ECLDCR). *Graefe's Archive for Clinical and Experimental Ophthalmology*. 2012;250(8):1249-1250.
24. Warren, J. F., Seiff, S. R., & Kavanagh, M. C. (2005). Long-term results of external dacryocystorhinostomy.
25. Dolman, P. J. (2003). Comparison of external dacryocystorhinostomy with nonlaser endonasal dacryocystorhinostomy. *Ophthalmology*, 110(1), 78-84.
26. Mourya, D., & Rijal, R. K. (2017). Transcanalicular laser-assisted dacryocystorhinostomy with diode laser. *Orbit*, 36(6), 370-374.
27. Henson, R. D., Henson Jr, R. G., Cruz Jr, H. L., & Camara, J. G. (2007). Use of the diode laser with intraoperative mitomycin C in endocanalicular laser dacryocystorhinostomy. *Ophthalmic Plastic & Reconstructive Surgery*, 23(2), 134-137.
28. Ajalloueyan, M., Fartookzadeh, M., & Parhizgar, H. (2007). Use of laser for dacryocystorhinostomy. *Archives of Otolaryngology-Head & Neck Surgery*, 133(4), 340-343.
29. Narioka, J., & Ohashi, Y. (2008). Transcanalicular-Endonasal Semiconductor Diode Laser-Assisted Revision Surgery for Failed External

- Dacryocystorhinostomy. *American journal of ophthalmology*, 146(1), 60-68.
30. Gupta, S. K., Kumar, A., Agarwal, S., & Pandey, P. (2012). Transcanalicular laser dacryocystorhinostomy using low energy 810 nm diode laser. *Oman journal of ophthalmology*, 5(3), 171.
 31. Ayintap, E., Buttanri, I. B., Sadigov, F., Serin, D., Ozsutcu, M., Umurhan Akkan, J. C., & Tuncer, K. (2014). Analysis of age as a possible prognostic factor for transcanalicular multidiode laser dacryocystorhinostomy. *Journal of ophthalmology*, 2014.
 32. Koch, K. R., Cursiefen, C., & Heindl, L. M. (2015). Transcanalicular laser dacryocystorhinostomy: one-year-experience in the treatment of acquired nasolacrimal duct obstructions. *Klinische Monatsblätter für Augenheilkunde*, 233(2), 182-186.
 33. Mor, J. M., Matthaei, M., Schruppf, H., Koch, K. R., Bölke, E., & Heindl, L. M. (2018). Transcanalicular laser dacryocystorhinostomy for acquired nasolacrimal duct obstruction: an audit of 104 patients. *European journal of medical research*, 23(1), 1-7.
 34. Taşkıran Çömez, A., Karadağ, O., Arıkan, S., Gencer, B., & Kara, S. (2014). Comparison of transcanalicular diode laser dacryocystorhinostomy and external dacryocystorhinostomy in patients with primary acquired nasolacrimal duct obstruction. *Lasers in surgery and medicine*, 46(4), 275-280.
 35. Derya, K., Demirel, S., Doganay, S., Orman, G., Cumurcu, T., & Gunduz, A. (2013). Endoscopic transcanalicular diode laser dacryocystorhinostomy: is it an alternative method to conventional external dacryocystorhinostomy?. *Ophthalmic Plastic & Reconstructive Surgery*, 29(1), 15-17.
 36. Balikoglu-Yilmaz, M., Yilmaz, T., Taskin, U., Taskapili, M., Akcay, M., Oktay, M. F., & Eren, S. (2015). Prospective comparison of 3 dacryocystorhinostomy surgeries: external versus endoscopic versus transcanalicular multidiode laser. *Ophthalmic Plastic & Reconstructive Surgery*, 31(1), 13-18.
 37. Akcay, E., Yuksel, N., & Ozen, U. (2016). Revision external dacryocystorhinostomy results after a failed dacryocystorhinostomy surgery. *Ophthalmology and therapy*, 5(1), 75-80.
 38. Henson, R. D., Cruz, H. L., Henson Jr, R. G., Ali, M. J., & Kakizaki, H. (2012). Postoperative application of mitomycin-C in endocanalicular laser dacryocystorhinostomy. *Ophthalmic Plastic & Reconstructive Surgery*, 28(3), 192-195.
 39. Basmak, H., Cakli, H., Sahin, A., Gursoy, H., Ozer, A., & Colak, E. (2011). What is the role of partial middle turbinectomy in endocanalicular laser-assisted endonasal dacryocystorhinostomy?. *American journal of rhinology & allergy*, 25(4), e160-e165.
 40. Onerci, M. (2002). Dacryocystorhinostomy. Diagnosis and treatment of nasolacrimal canal obstructions. *Rhinology*, 40(2), 49-65.
 41. Mak, S. T., Io, I. Y. F., & Wong, A. C. M. (2013). Prognostic factors for outcome of endoscopic dacryocystorhinostomy in patients with primary acquired nasolacrimal duct obstruction. *Graefe's Archive for Clinical and Experimental Ophthalmology*, 251(5), 1361-1367.
 42. Smirnov, G., Pirinen, R., Tuomilehto, H., Seppä, J., Teräsvirta, M., Uusitalo, H., ... & Kaarniranta, K. (2011). Strong expression of HSP47 in metaplastic nasal mucosa may predict a poor outcome after primary endoscopic dacryocystorhinostomy: a prospective study. *Acta ophthalmologica*, 89(2), e132-e136.
 43. Linberg, J. V., Anderson, R. L., Bumsted, R. M., & Barreras, R. (1982). Study of intranasal ostium external dacryocystorhinostomy. *Archives of Ophthalmology*, 100(11), 1758-1762.
 44. Ezra, E., Restori, M., Mannor, G. E., & Rose, G. E. (1998). Ultrasonic assessment of rhinostomy size following external dacryocystorhinostomy. *British journal of ophthalmology*, 82(7), 786-789.
 45. Limbu, B., Sim, B., Shrestha, M. K., Tabin, G., & Saiju, R. (2020). Results of early versus standard silicone stent removal following external dacryocystorhinostomy under local anesthesia. *Asian Journal of Ophthalmology*, 17(3), 263-272.