

Isolation of an Anti-Thiamine Compound from *Bergenia ciliata* leaves of Sikkim Himalaya

Tanaya Ghosh¹, Prasanta Kumar Mitra^{1*}

¹Department of Medical Biotechnology, Sikkim Manipal University, Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim, India

DOI: [10.36348/sjbr.2022.v07i01.006](https://doi.org/10.36348/sjbr.2022.v07i01.006)

Received: 14.12.2021 | Accepted: 23.01.2022 | Published: 29.01.2022

*Corresponding author: Prasanta Kumar Mitra

Department of Medical Biotechnology, Sikkim Manipal University, Sikkim Manipal Institute of Medical Sciences, Gangtok, Sikkim, India

Abstract

Recently we have shown that *Bergenia ciliate* (*B. ciliate*, Family- Saxifragaceae) leaves of the period July – August had maximum in vitro anti thiamine effect. Aim of the present work was, to isolate anti thiamine compound from the leaves of *B. ciliate*. From the local market *B. ciliate* leaves were collected during July – August and identified by the taxonomist. Solvent extraction and acid hydrolysis were done followed by solvent treatment, chromatographic experiments and crystallization. A compound was crystallized. In vitro anti thiamine activity of the isolated compound was checked by standard method. Results revealed that 100 mg of the isolated compound could inactivate 29.5 mg of thiamine hydrochloride in in vitro experiment.

Keywords: *Bergenia ciliate* leaves, isolated compound, anti thiamine activity.

Copyright © 2022 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Anti thiamine activity of the plant was established as early as 1944 when Bhagvat *et al.* demonstrated anti thiamine activity in *Linum usitatissimum*. Authors claimed that the anti-thiamine factor present in the plant was thermostable and largely dialyzable [1]. In 1946 anti thiamine activity of *Pteris aquilina* L. was confirmed in experimental animals, bacteria and human [2]. Haag *et al.*, showed that the anti thiamine factor of bracken fern was very stable to heat, it was water soluble but was not soluble in petroleum ether, acetone or ethanol. Inactivation of thiamine by bracken fern extract was produced rapidly [3]. Fabriani *et al.*, confirmed anti thiamine activity of *Pteris aquilina* [4]. In 1952 Henderson *et al.*, reported that the plant *Pteris aquilina* contains thiaminase which causes thiamine deficiency in horses [5]. In 1956 Sakamoto and coworkers isolated a thiamine-decomposing thermostable factor from sweet potato leaves which was characterized as isoquercitrin [6].

Chaudhuri confirmed the presence of a heat stable thiamine inactivating-factor in different varieties of rice and rice bran [7]. In 1964 Kundig & Somogyi found that aqueous extract of parsley contained an anti thiamine factor which was unaffected by cooking or by

contact with synthetic gastric juice [8]. Rattanapanone and coworkers showed anti thiamine activity of water extract of *Marsilea crenata*. They noted that the plant has enzyme thiaminase and a thermostable factor responsible for anti thiamine activity. These workers also confirmed anti thiamine activity of several other plants including *Athyrium esculentum*, an edible fern (Family-Athyriaceae) found throughout Asia and Oceania [9]. Kositawattanukul and his colleagues found that tea plant has anti thiamine activity which is protected by ascorbic acid (vitamin C) both at acidic and neutral pH [10]. We have also noted *in vitro* anti thiamine activity (70% inhibition of thiamine was noted in 120 minutes) of *Abrus precatorius* L. leaves [11].

Recently we have confirmed *in vitro* anti thiamine activity of *B. ciliate* leaves (results are under communication). We report here isolation of the anti thiamine compound from *B. ciliate* leaves.

MATERIAL AND METHODS

Collection of plant material

Fresh and healthy leaves of *B. ciliata* were collected from the local market during July – August of Gangtok, Sikkim (we have seen earlier that *B. ciliata* leaves of July – August has maximum anti thiamine

activity) & identified by the taxonomist. Voucher specimen (No. SM-MB-010/21-5) was kept in the department of Medical Biotechnology, Sikkim Manipal University for future references.



Bergenia ciliata leaves

Preparation of leaves for Anti thiamine activity

Leaves of *B. ciliata* were shed dried and powdered. This powder was used to isolate the anti thiamine compound.

In vitro anti-thiamine activity

The anti thiamine activity was determined by estimating the residual thiamine present in a system containing known amount of thiamine hydrochloride and test material collected from *B. ciliata* leaves by the method of Bhattacharya and Choudhuri [12].


Main steps were: an intimate mixture of thiamine hydrochloride (100 mg) and isolated compound from *B. ciliata* leaves (100 mg) was incubated at 30 degree centigrade for 1 hour in 10 ml M/15 phosphate buffer at pH 6.5. It was then filtered. 2 ml of this filtrate was taken and residual thiamine hydrochloride was estimated by thiochrome method described by Harris and Wang [13]. In short, to 2ml of the filtrate 0.1ml potassium ferricyanide (2.5g/l) and 0.25 ml of sodium hydroxide (150g/l) were added. The solution was mixed thoroughly. 2 ml isobutanol was then added to it. The solution was shaken for 1 minute. Fluorescence of the supernatant was noted by a fluorimeter at 435 nm using excitation at 365 nm. Tubes for standard thiamine solution (400 µg/l) and for blank were run simultaneously.

Reagents

All chemicals used in this study were purchased from Sigma Chemical Company, Mumbai. Chemicals were of analytical grade with high purity

Isolation of alpha amylase inhibitor from *B. ciliata* Leaves

Applying principles of standard isolation procedures of chemical compounds from plant sources [14, 15], isolation was done by the following scheme.



RESULTS

Isolation of compound

One compound was isolated from leaves of *B. ciliata*.

Anti thiamine activity of the isolated compound

Results are summarized in Table-1.

Table-1: Showing in vitro anti thiamine effect of the isolated compound from *B. ciliata* leaves

Group	Residual thiamine (mg)	Inhibition (%)
Control (Thiamine hydrochloride)	100.0	--
Thiamine hydrochloride (100 mg) + Isolated compound from <i>B. ciliata</i> leaves (100 mg).	70.5 ± 2.5*	29.5

Values were mean ± SEM of ten sets of experiment. *p < 0.05, when compared to control.

In *in vitro* experiment 100 mg of the isolated compound could destroy 29.5 mg thiamine. Initially amount of thiamine was 100 mg. After 1h incubation with 100 mg of the isolated compound, amount of thiamine came down to 70.5 ± 2.5 . Result was statistically significant. Percentage of thiamine destruction was 29.5%.

DISCUSSION

The concept of anti-thiamine factor was introduced in literature by Green [16]. Evans *et al.*, (1942) classified anti-thiamine compounds broadly into two categories, namely synthetic and natural. Synthetic anti-thiamine compounds are structural analogues or

Antimetabolites e.g. pyriothiamine type, oxythiamine type, amproleum type, deoxy and ethyl deoxy thiamine, O-benzoyl thiamine and its derivatives, butyl thiamine, phenyl triazinothiamine, imidazolethiamine and benzoyl imidazole thiamine etc.; while natural anti thiamine compounds are non-structural analogues and mostly present in different food-stuffs, plants etc. Natural anti thiamine compounds are further classified into two groups *viz.* large molecule natural anti-thiamine compounds e.g. thiaminase I and thiaminase II, mainly isolated from raw fishes and small molecule natural anti-thiamine compounds e.g. caffeic acid, chlorogenic acid, methyl sinapate, 3-5 dimethoxy salicylic acid etc [17].

Plants are known having anti thiamine activity and many anti thiamine compounds have been isolated from plants. Reddi in 1950 isolated an active principle from mustard seed which destroyed thiamine. The factor is soluble in water and alcohol, sparingly soluble in chloroform but insoluble in ether and acetone. It was found out that 65% of the added thiamine was destroyed by the anti thiamine factor even within the first minute. The destruction of thiamine was independent of pH 2.5 and 7.5. The reaction was not enzymatic [18].

In 1971 Hilker and co-workers made some extensive studies on thiamine-inactivating factors present in various types of tea such as jasmine, black tea, etc. Authors demonstrated that thiamine-inactivating factors of tea appeared to be related to the tannin content [19]. Further, it was demonstrated that ortho-catechol such as methylsinapate, found in rapeseed, has antithiamine activity [20].

Bhattacharya and coworkers in 1974 confirmed that mustard seed contains methyl ester of 3, 5-dimethoxy-4-hydroxy cinnamic acid (methyl sinapate) which acts as anti thiamine factor. It was found out that 1 mg of methyl sinapate inactivated 45 µg of thiamine hydrochloride at pH 6.5 at 30°C. Under similar conditions 1 mg of caffeic acid (3, 4-dihydroxy cinnamic acid) inactivated 135 µg of thiamine

hydrochloride. Maximum inactivation of thiamine by methyl sinapate was recorded within 30 min [12].

The anti thiamine factor, isolated from cotton seed, was a light yellow amorphous substance. The anti thiamine factor was characterized as 3, 5-dimethoxy salicylic acid [21].

In 1980 Wills *et al.*, studied anti thiamine activity of tea and isolated a fraction from tea which had very strong anti thiamine effect [22].

In the present study we have isolated an anti thiamine compound from *B. ciliata* leaves, 100 mg of which could inactivate 29.5 mg of thiamine hydrochloride. The isolated compound should need characterization. Work in this direction is now in progress in our laboratory.

CONCLUSION

Thiamine is the coenzyme of both pyruvate dehydrogenase and transketolase enzymes. Activity of these enzymes will decrease in case of presence of anti thiamine compound in edible stuff. Due to this, energy production as well as synthesis of nucleic acid will be hampered. Anti thiamine activity of *B. ciliate* leaves, therefore, should be known to the patients who are taking *B. ciliate* leaves to get rid of various ailments.

Conflict of interest: The authors declare that they have no conflict of interest.

REFERENCES

1. Bhagvat, K., & Devi, P. (1944). Inactivation of thiamine by certain foodstuffs and oil seeds. Part-II. *Indian Journal of Medical Research*, 32, 139-144.
2. Weswig, P. H., Freed, A. M., & Haag, J. R., (1946). Antithiamine activity of plant materials. *Journal of Biological Chemistry*, 165(2), 73-738.
3. Haag, J. R., Weswig, P. H., & Freed, A. M. (1947). Antithiamine activity of bracken fern. *Federation Proceedings*, 6, 408.
4. Fabriani, J., Fratoni, A., & Spadoni, M. A. (1948). Azione biologica della tiaminase nella modificazione in essa provocata da un fattore contenuto nelle foglie di felce di macchia (*Pteris aquilina*). *Q Nutr*, 10, 295.
5. Henderson, J. A., Evans, E. V., & McIntosh, R. A. (1952). The antithiamine action to *Equisetum*. *Journal of the American Veterinary Medical Association*, 120, 375-378.
6. Sakamoto, S. and Fujita, A. (1956). studies on thiamine-decomposing thermostable factors. II. Isolation of thiamine-decomposing thermostable factor from sweet potato leaves. *The Journal of Vitaminology*, 2(1), 39-43.
7. Chaudhuri, D. K. (1962). Antithiamine factor of rice-bran, *Science and Culture*, 28, 384.

8. Kundig, H., & Somogyi, J. C. (1964). Antithiamin wirkstoffe in pflanzlichen Nahrungsmitteln. *Int J Vitaminforsch*, 34, 135.
9. Rattanapanone, V., Sanpitak, N., & Phornphiboul, B. (1971). A study of antithiamine factor in raw vegetables of northern part of Thailand. *Chiang Mai Medical Bulletin*, 10(1), 17-23.
10. Kositawattanukul, T. (1977). Chemical interactions between thiamine and tannic acid. II. Separation of products. *The American Journal of Clinical Nutrition*, 30(10), 1686-1691.
11. Mitra, P. K. (2014). Effect of solvent, temperature, pH and time on extraction process of anti thiamine factor present in leaves of *Abrus precatorius* linn. *World Journal of Pharmaceutical Research*, 3(3), 4816-4824.
12. Bhattacharya, J., & Chaudhury, D. K. (1974). Antithiamine factor present in *Brassica juncea*. *Biochem Biophys. Acta*, 343, 211-220.
13. Harris, L. J., & Wang, Y. L. (1941). A new method for estimation of residual thiamine. *Biochem J*, 35, 1050-1058.
14. Cannell, R. J. P. (1998). Natural Products Isolation, New Jersey, Human Press Inc. 165-208.
15. Li, H. B., Jiang, Y., & Chen, F. (2004). Separation methods used for *Scutellaria baicalensis* active components. *J. Chromatography*, 8(12), 277-290.
16. Green, R. G. (1936). Studies on Chastekparalysis, *Minn Wildl Dis Inv*, 2, 106-113.
17. Evans, C. A., Carison, W. E., & Green, R. G. (1942). Chastek paralysis – Role of thiamine. *American Journal of Pathology*, 18, 79.
18. Reddi, K. K. (1950). Studies on antithiamine factor in mustard (*Brassica juncea*), *J Biol Sci*, 17(1), 39-44.
19. Hilker, D. M. (1971). Antithiamine effects of tea: temperature and pH dependence, *Nutrition Reports International*, 4, 223.
20. Noda, M., & Matszjimoto, M. (1971). Sinapic acid and methyl sinapate in rapeseed lipids. *Biochimica biophysica acta*, 231, 131-133.
21. Sarkar, L., & Chaudhuri, D. K. (1976). Antithiamine factor in cotton seed (*Bombex malabericum*)-its isolation and characterization. *International Journal of Vitamin and Nutrition Research*, 46(4), 417-421.
22. Wills, R. B. H., & Mcbrien, K. J. (1980). Antithiamin activity of tea fractions, *Foodchemistry*, 6(2), 111-114.