

Review Article

Influence of Impaired Masticatory Function on the Nutrition of Completely Edentulous Patients

Suryakant C. Deogade¹, Dinesh Naitam², Sushant M. Patil³,

¹Professor, Department of Prosthodontics and Crown & Bridge, Hitkarini Dental College & Hospital, Jabalpur, Madhya Pradesh, India

²Asso. Professor, Department of Dentistry, Government Medical College, Akola, Maharashtra, India

³Dental Surgeon, Department of Prosthodontics and Crown & Bridge, Government Dental College and Hospital, Nagpur, Maharashtra, India

*Corresponding Author:

Dr.Suryakant C. Deogade

Email: dr_deogade@yahoo.co.in

Abstract: Individuals suffering from impaired masticatory function (MF) may adapt food consistency to their existing dental status or rely on the digestive system to compensate for the lack of oral preparation of food. These circumstances further leads to deficient nutrient intake or increase the likelihood of digestive diseases and decrease gut absorption. Dysfunction of masticatory efficiency thus may be detrimental to general health. This article reviews evidence of the effects of masticatory deficiency on nutrition. It is difficult to draw conclusions from many of the reviewed studies due to issues related to study design, confounding variables, and the subjective nature of the measurements. The reviewed evidences supporting an association between MF and deficient dietary intake often are based on relatively weak correlations and cannot confer a causal relationship between the reviewed variables.

Keywords: mastication, nutrition, digestion, diet

INTRODUCTION

Loss of tooth following its removal from mouth leads to a frequent condition of deficient masticatory function (MF). Most common causes of tooth loss include caries and periodontal disease, which may result in total tooth loss causing a complete edentulism in patients. Manly and Braley [1] have reported that the prevalence of edentulism in persons aged 65 to 75 years is 16.9% in France, 24.8% in Germany, and 26% to 31% in the United States. They found that the masticatory performance and efficiency of a denture wearer is considerably less compared to that of a fully dentate individual.

Diminished MF can result in functional deficiencies of the tongue, the oral mucosa, the muscles of mastication, the salivary glands, and the nervous system. The possible effects of this impairment of MF on the general health of an individual are not fully understood and remain unclear. Individuals with impaired MF may adapt their food choices or swallow coarse particles that make the problem a digestive one. This type of behavior induce imbalance in dietary intake or may result in decreased bioavailability of nutrients and gastrointestinal disturbances in such people. In both situations, the impaired dietary or nutrient intake may increase the incidence of nutrition-deficient conditions or disease [2]. This article reviews

the impact of impaired MF on nutrition and health of completely edentulous individuals.

EVALUATION OF MASTICATORY FUNCTION

Several authors [3,4] designed the different methods for evaluation of masticatory efficiency, ability, and performance. These workers suggested five different categories. The first category involves self-assessment of MF by means of scales and questionnaires. These methods are used in epidemiological surveys, where the participating individuals are asked to score their ability to chew foods into good, fairly good, or poor or to rate foods into easy, fairly easy, difficult, or very difficult to chew. Slagter and colleagues[4] advocated that this method provides very optimistic results compared to a practitioner's evaluation and requires a large sample population for survey. The second method measures the individual's ability to comminute food into smaller particles, where the chewed test food is retrieved and analyzed after mastication for a set number of strokes or set time.³⁻¹³The retrieved test food is then screened through sieves with a range of meshes, from which the masticatory index is calculated. Some authors also suggested the swallowing threshold test index, where the individuals are instructed to chew a standard portion of test food in as many strokes as required until they are ready to swallow it. This chewed portion of test food is then expectorated and analyze through sieving [5, 12,

14, 15]. The swallowing threshold test index is considered as an objective method as time and the number of chewing strokes are not limited and involves the natural conditions of bolus preparation. Heath [16] suggested the third method that measures the amount of sugar extracted from chewing gum. The fourth method encounters the different techniques which include bite force measurement, electromyography, kinematics, and video recording [16-20]. However, these methods demands expensive equipment and special training and need to perform in a large-scale epidemiological survey increasing its difficulties for its implication. The fifth method considers certain anatomic criteria, where each tooth is assigned a numerical coefficient according to its presumed functional importance [11, 21, 22]. These anatomical based methods encounters the teeth and neglect the role of other masticatory organs such as salivary glands, tongue, lips, cheeks and muscles of mastication.

FACTORS RESPONSIBLE FOR IMPAIRED MASTICATORY EFFICIENCY

Several authors [23-27] reported that the tooth loss leads to impairment of masticatory function. Gunne⁶ studied the effect of removable partial dentures on mastication and dietary intake. He suggested that the replacement of missing teeth with removable dentures can improve mastication. Contrary to this, other authors [6, 7, 14, 28, 29] believed that such replacement cannot recover the efficiency of a dentate state. Kubota and colleagues [30] supported the degenerative changes of primary neurons following tooth extraction as one of the reason behind impaired MF. Gobel and Binck [31] also reported the similar findings in their animal studies. Appenteng and colleagues [32] investigated the intraoral mechanoreceptor activity during jaw movement in the anesthetized rabbit and found that the completed dentures often cover large areas of the oral mucosa and block part of its sensory activity. Several studies [33-36] have reported an impaired masticatory efficiency due to dysfunction of tongue motor skills and lack of tonicity of muscles involved in masticatory movements. Watanabe and colleagues³⁷ stressed the role of the tongue in the collection of sensory information and preparing the food bolus ready for swallowing. Liedberg and Owall [34] investigated the masticatory ability in experimentally induced xerostomia and found that such condition may negatively influence the masticatory process by making it impossible for individuals to convert food into bolus form before swallowing.

CORRELATION BETWEEN IMPAIRED MF AND DEFICIENT DIETARY INTAKE

The correlation between MF and nutrition has not been established. Experimental and longitudinal studies may not reveal this relationship due to various reasons. However, numerous authors [23, 26, 29, 38-49] have tried to correlate the impaired MF with that of inadequate food selection in edentulous people. In some

western countries, the edentulous populations have a wide choice of foods. The individuals suffering from impaired MF may adapt food consistency to their existing dentate condition, which may result in an increased consumption of soft foods and a restricted consumption of hard-to-chew foods [45-49]. Several authors [45, 48, 49, 50, 51] categorized the food choice opted by an edentulous subject. Individuals with impaired MF may choose to eat industrially processed instead of natural foods. Fillion and Henry [52] reviewed the various forms of industrial food processing and concluded that in patients with impaired MF, industrial foods may favor the absorption of substantial amounts of fat and increase the eater's level of cholesterol and saturated fatty acids. In other studies [38, 39, 44, 47, 50, 53], the individuals were found to avoid hard-to-chew natural foods such as crunchy foods (raw vegetables and fresh fruits), stringy foods (meat), and dry fruits (bread or bagels) from their diet.

Renaud and colleagues [47] discussed dietary problems before and after rehabilitation of the MF in edentulous subjects. They observed that soft fruits (raspberries, strawberries, grapes, and tomatoes) may be avoided by denture wearers because their seeds often lodge under prostheses. Some authors [54, 55] reported that denture wearers may choose home processing to soften foods, which may alter the nutritional value of foods due to long cooking times. This causes degradation of essential micronutrients such as vitamin C, thiamin, and folates and reduces the expected bioavailability of these nutrients in the ingested food. Home processing of foods may also produce trans-fatty acids [52]. Krall and colleagues [51] compared the dietary intake in complete denture wearers and dentate subjects and found that the protein and dietary fiber intake in former group was 15% and 24% lower, respectively, than that of later group. They also observed a decrease of 20%, 35%, and 27% in the intake levels of folic acid, carotene, and iron, respectively. Other authors [50, 56, 57] reported a significant decrease in the intake of vitamins C, A, B1, B2, and B12, folic acid, and essential minerals such as calcium and iron in complete denture wearers. Joshipura and colleagues [50] surveyed more than 49,000 dentate and edentulous male health care professionals and observed that the mean differences in the intake of 10 different nutrients ranged from 2% to 13%. In partially edentulous subjects the decrease in nutrient intake was proportional to the number of missing teeth.

Hamada and colleagues [58] carried a randomized clinical trial comparing the efficacy of mandibular implant-supported overdentures and conventional dentures in diabetic patients and failed to demonstrate substantial changes in patient nutrient intake after an improvement in dental status. Authors related the bad dietary habits as the causative factor rather than the poor dental status. Horwath [41]

suggested that some studies assessed MF on the basis of questionnaire which could not reflect the reality of dental arch efficiency. Overall a weak correlation between MF and dietary intake has been reported and needs the future sound scientific studies to reveal any potential relationship.

IMPAIRED MF AND RESULTING NUTRITIONAL CONSEQUENCES

Some workers [33, 42] carried experiments in animals like rats and pigs and observed that mastication exhibits a moderate influence on the digestion of foods. Kapur and Okubo [59] reported the similar findings in rats and claimed a more evident influence of impaired MF on the digestion of foods in old than in young rats. Geissler [42] found that minced meat is at best poorly digested by dogs but meat is completely digested when swallowed in large lumps. These findings suggest that the conclusions drawn from animal and human studies varied. Hence, species differences should be considered in this area of research. Prinz and Lucas [15] experimented mixtures of solid and liquid food in various proportions in human mastication and reported that swallowing threshold is related to both particle size and degree of moistening. Several authors [60, 61] studied the absorption of whole peanuts, peanut oil, peanut butter and cereal food and observed that insufficient mastication of these foods remained intact in the feces. Rodriguez-Olleros and Rey [62] reported gastritis and ulcers in subjects with impaired MF. Hedde and colleagues [63] carried animal (swine) experiments and concluded that the results appeared to be dependent on the species. Swines do not exhibit the problems similar to those of humans. Brodeur and colleagues [44] suggested the correlation between masticatory efficiency and gastrointestinal disturbances, as expressed by the presence of symptoms such as diarrhea and constipation is mostly seen in the elderly.

EFFECT OF IMPAIRED DIETARY INTAKE ON VARIOUS DISEASE RISKS

Several studies [64-69] have reported correlation between poor oral health and general health. Many micronutrients and macronutrients provide protection against various types of diseases, and the excessive consumption of some nutrients may be harmful to health. Several epidemiologic investigations [70-78] conducted in various countries have found that intake of vegetables and fruits (raw carrots, pears, and apples) can lower the prevalence of cancer at many sites. Block and colleagues [74] reviewed the epidemiological evidence of fruits and vegetables in prevention of cancer involving lung, colon, cervix, esophagus, oral cavity, stomach, bladder, pancreas or ovary. They reviewed around 156 articles, out of which 128 concluded that fruit and vegetable consumption had a significantly protective effect. Scheppach and colleagues [77] suggested the existence of many potential anticarcinogenic agents in vegetables and fruits. The authors suggested the possible mechanism

behind anticarcinogenic potential occurring via an inhibition of nitrosamine formation, antioxidant effects, provision of substrates for formation of antineoplastic agents, and dilution and binding of carcinogens. The dietary fiber decreases the colorectal cancer mortality by binding bile acids, lowering acidity, and increasing fecal bulk [78]. Schneeman [79] reported the active role of dietary fiber in facilitating gastrointestinal transit, lowering plasma cholesterol levels, reducing the glycemic response to carbohydrate-containing meals. Jenkins and colleagues [80] has recommended the consumption of dietary fiber for the prevention of Crohn's disease, constipation, irritable bowel syndrome, diverticular disease, and gallstones. Werler and colleagues [81] reported that low consumption of vitamin C has been inversely associated with cataracts and slow healing after injury. A daily periconceptional consumption of folic acid by women may lower the risk of neonatal neural tube defects. Several authors [82-84] reported the potential role of saturated fatty acids in developing atherosclerosis. Increased cholesterol and saturated fatty acid consumption increases the risk of cardiovascular diseases especially myocardial infarction. Some studies [85, 86] revealed the role of a fatty diet in increasing the risk of hypertension, cardiovascular diseases and noninsulin-dependent diabetes. Thus, impaired MF could be considered as indirect risk factor for the etiology of cardiovascular diseases.

CONCLUSION

A compromised oral health causes an impairment of MF, which further may lead to inadequate choice of foods altering nutrient consumption. This suggests that a dysfunction in masticatory efficiency is correlated with the disturbances into general health of an edentulous individual. A significant correlation between a poor oral health and some gastrointestinal diseases has been established. Public health services must be given stress on the need for preserving natural teeth to enable adequate nutritional consumption, and dental practitioners should be aware of various nutritional risks associated with denture wearing.

REFERENCES

1. Manly, R. S., & Braley, L. C. (1950). Masticatory performance and efficiency. *Journal of Dental Research*, (29), 448-62.
2. N'Gom, P. I., & Woda, A. (2002). Influence of impaired mastication on nutrition. *The Journal of prosthetic dentistry*, 87(6), 667-673.
3. Gunne, H. S. J., & Wall, A. K. (1985). The effect of new complete dentures on mastication and dietary intake. *Acta Odontologica*, 43(5), 257-268.
4. Slagter, A. P., Olthoff, L. W., Bosnian, F., & Steen, W. H. (1992). Masticatory ability, denture quality, and oral conditions in

- edentulous subjects. *The Journal of prosthetic dentistry*, 68(2), 299-307.
5. Feldman, R. S., Kapur, K. K., Alman, J. E., & Chauncey, H. H. (1980). Aging and Mastication: Changes in Performance and in the Swallowing Threshold with Natural Dentition†. *Journal of the American Geriatrics Society*, 28(3), 97-103.
 6. Gunne, H. S. J. (1985). The effect of removable partial dentures on mastication and dietary intake. *Acta Odontologica*, 43(5), 269-278.
 7. Gunne, H. S. (1985). Masticatory efficiency and dental state. A comparison between two methods. *Acta Odontologica Scandinavica*, 43(3), 139-146.
 8. Akeel, R., Nilner, M., & Nilner, K. (1991). Masticatory efficiency in individuals with natural dentition. *Swedish dental journal*, 16(5), 191-198.
 9. Slagter, A. P., Olthoff, L. W., Steen, W. H. A., & Bosman, F. (1992). Comminution of food by complete-denture wearers. *Journal of dental research*, 71(2), 380-386.
 10. Slagter, A. P., Bosman, F., Van Der Glas, H. W., & Van Der Bilt, A. (1993). Human jaw-elevator muscle activity and food comminution in the dentate and edentulous state. *Archives of oral biology*, 38(3), 195-205.
 11. Julien, K. C., Buschang, P. H., Throckmorton, G. S., & Dechow, P. C. (1996). Normal masticatory performance in young adults and children. *Archives of oral biology*, 41(1), 69-75.
 12. Garrett, N. R., Perez, P., Elbert, C., & Kapur, K. K. (1996). Effects of improvements of poorly fitting dentures and new dentures on masticatory performance. *The Journal of prosthetic dentistry*, 75(3), 269-275.
 13. Kapur, K., Soman, S., & Yurkstas, A. (1964). Test foods for measuring masticatory performance of denture wearers. *The Journal of Prosthetic Dentistry*, 14(3), 483-491.
 14. Kapur, K. K., & Soman, S. D. (1964). Masticatory performance and efficiency in denture wearers. *The Journal of Prosthetic Dentistry*, 14(4), 687-694.
 15. Prinz, J. F., & Lucas, P. W. (1995). Swallow thresholds in human mastication. *Archives of oral biology*, 40(5), 401-403.
 16. Heath, M. R. (1982). The effect of maximum biting force and bone loss upon masticatory function and dietary selection of the elderly. *International dental journal*, 32(4), 345-356.
 17. Diaz-Tay, J., Jayasinghe, N., Lucas, P. W., McCallum, J. C., & Jones, J. T. (1991). Association between surface electromyography of human jaw-closing muscle and quantified food breakdown. *Archives of oral biology*, 36(12), 893-898.
 18. Lassauzay, C., Peyron, M. A., Albuissou, E., Dransfield, E., & Woda, A. (2000). Variability of the masticatory process during chewing of elastic model foods. *European journal of oral sciences*, 108(6), 484-492.
 19. Peyron, M. A., Maskawi, K., Woda, A., Tanguay, R., & Lund, J. P. (1997). Effects of food texture and sample thickness on mandibular movement and hardness assessment during biting in man. *Journal of Dental Research*, 76(3), 789-795.
 20. Schwartz, J. L., Niman, C. W., & Gisel, E. G. (1984). Chewing cycles in 4-and 5-year-old normal children: an index of eating efficacy. *American Journal of Occupational Therapy*, 38(3), 171-175.
 21. Miura, H., Araki, Y., Hirai, T., Isogai, E., Hirose, K., & Umenai, T. (1998). Evaluation of chewing activity in the elderly person. *Journal of oral rehabilitation*, 25, 190-193.
 22. Wilding, R. J. C. (1993). The association between chewing efficiency and occlusal contact area in man. *Archives of oral biology*, 38(7), 589-596.
 23. Chauncey, H. H., Muench, M. E., Kapur, K. K., & Wayler, A. H. (1984). The effect of the loss of teeth on diet and nutrition. *International Dental Journal*, 34(2), 98-104.
 24. Tsuga, K., Carlsson, G. E., Österberg, T., & Karlsson, S. (1998). Self-assessed masticatory ability in relation to maximal bite force and dental state in 80-year-old subjects. *Journal of oral rehabilitation*, 25(2), 110-116.
 25. Agerberg, G., & Carlsson, G. E. (1981). Chewing ability in relation to dental and general health: analyses of data obtained from a questionnaire. *Acta Odontologica*, 39(3), 147-153.
 26. Wayler, A. H., & Chauncey, H. H. (1983). Impact of complete dentures and impaired natural dentition on masticatory performance and food choice in healthy aging men. *The Journal of prosthetic dentistry*, 49(3), 427-433.
 27. Miura, H., Araki, Y., & Umenai, T. (1997). Chewing activity and activities of daily living in the elderly. *Journal of oral rehabilitation*, 24(6), 457-460.
 28. CHONG-SHAN, S. H. I., GUAN, O., & TIAN-WEN, G. U. O. (1990). Comparison of food particle distribution masticated by subjects wearing complete dentures and with natural teeth. *Journal of oral rehabilitation*, 17(6), 611-615.
 29. Ettinger, R. L. (1998). Changing dietary patterns with changing dentition: how do people cope?. *Special Care in Dentistry*, 18(1), 33-39.
 30. Kubota, K., Nagae, K., Shibana, S., Hosaka, K., Iseki, H., Odagiri, N., ... & Narita, N. (1987). Degenerative changes of primary

- neurons following tooth extraction. *Anatomischer Anzeiger*, 166(1-5), 133-139.
31. Gobel, S., & Binck, J. M. (1977). Degenerative changes in primary trigeminal axons and in neurons in nucleus caudalis following tooth pulp extirpations in the cat. *Brain Research*, 132(2), 347-354.
 32. Appenteng, K., Lund, J. P., & Seguin, J. J. (1982). Intraoral mechanoreceptor activity during jaw movement in the anesthetized rabbit. *Journal of Neurophysiology*, 48(1), 27-37.
 33. Carlsson, G. E. (1974). Bite force and chewing efficiency.
 34. Liedberg, B., & Öwall, B. (1991). Masticatory ability in experimentally induced xerostomia. *Dysphagia*, 6(4), 211-213.
 35. Koshino, H., Hirai, T., Ishijima, T., & Ikeda, Y. (1997). Tongue motor skills and masticatory performance in adult dentates, elderly dentates, and complete denture wearers. *The journal of prosthetic dentistry*, 77(2), 147-152.
 36. Hennequin, M., Faulks, D., Veyrone, J. L., & Bourdiol, P. (1999). Significance of oral health in persons with Down syndrome: a literature review. *Developmental Medicine & Child Neurology*, 41(04), 275-283.
 37. Watanabe, S., Ohnishi, M., Imai, K., Kawano, E., & Igarashi, S. (1995). Estimation of the total saliva volume produced per day in five-year-old children. *Archives of oral biology*, 40(8), 781-782.
 38. Hildebrandt, G. H., Dominguez, B. L., Schork, M. A., & Loesche, W. J. (1997). Functional units, chewing, swallowing, and food avoidance among the elderly. *The Journal of prosthetic dentistry*, 77(6), 588-595.
 39. Baxter, J. C. (1984). The nutritional intake of geriatric patients with varied dentitions. *The Journal of prosthetic dentistry*, 51(2), 164-168.
 40. Sebring, N. G., Guckes, A. D., Li, S. H., & McCarthy, G. R. (1995). Nutritional adequacy of reported intake of edentulous subjects treated with new conventional or implant-supported mandibular dentures. *The Journal of prosthetic dentistry*, 74(4), 358-363.
 41. Horwath, C. C. (1990). Chewing difficulty and dietary intake in the elderly. *Journal of Nutrition for the Elderly*, 9(2), 17-24.
 42. Geissler, C. A., & Bates, J. F. (1984). The nutritional effects of tooth loss. *The American journal of clinical nutrition*, 39(3), 478-489.
 43. Shatenstein, B. (1986). Tooth loss, mastication, and nutrition: an overview. *Journal of Ethnopharmacology*, 17(3), 305.
 44. Brodeur, J. M., Laurin, D., Vallee, R., & Lachapelle, D. (1993). Nutrient intake and gastrointestinal disorders related to masticatory performance in the edentulous elderly. *The Journal of prosthetic dentistry*, 70(5), 468-473.
 45. Norlén, P., Steen, B., Birkhed, D., & Björn, A. L. (1993). On the relations between dietary habits, nutrients, and oral health in women at the age of retirement. *Acta Odontologica*, 51(5), 277-284.
 46. Ernest, S. L. (1993). Dietary intake, food preferences, stimulated salivary flow rate, and masticatory ability in older adults with complete dentitions. *Special Care in Dentistry*, 13(3), 102-106.
 47. Renaud, M., Mercier, P., & Gelinas, M. D. (1981). Dietary problems before treatment and after rehabilitation of the masticatory function. *Journal of the Canadian Dietetic Association*.
 48. Laurin, D., Brodeur, J. M., Bourdages, J., Vallee, R., & Lachapelle, D. (1994). Fibre intake in elderly individuals with poor masticatory performance. *Journal (Canadian Dental Association)*, 60(5), 443-6.
 49. Greksa, L. P., Parraga, I. M., & Clark, C. A. (1995). The dietary adequacy of edentulous older adults. *The Journal of prosthetic dentistry*, 73(2), 142-145.
 50. Joshipura, K. J., WILLET, W. C., & DOUGLASS, C. W. (1996). The impact of edentulousness on food and nutrient intake. *The Journal of the American Dental Association*, 127(4), 459-467.
 51. Krall, E., Hayes, C., & Garcia, R. (1998). How dentition status and masticatory function affect nutrient intake. *The Journal of the American Dental Association*, 129(9), 1261-1269.
 52. Fillion, L., & Henry, C. J. K. (1998). Nutrient losses and gains during frying: a review. *International journal of food sciences and nutrition*, 49(2), 157-168.
 53. Garcia, R. I., Perlmutter, L. C., & Chauncey, H. H. (1989). Effects of dentition status and personality on masticatory performance and food acceptability. *Dysphagia*, 4(2), 121-126.
 54. Yadav, S. K., & Sehgal, S. (1995). Effect of home processing on ascorbic acid and β -carotene content of spinach (*Spinacia oleracea*) and amaranth (*Amaranthus tricolor*) leaves. *Plant foods for human nutrition*, 47(2), 125-131.
 55. Severi, S., Bedogni, G., Zoboli, G. P., Manzieri, A. M., Poli, M., Gatti, G., & Battistini, N. (1998). Effects of home-based food preparation practices on the micronutrient content of foods. *European journal of cancer prevention*, 7(4), 331-336.
 56. Hamasha, A. A. H., Hand, J. S., & Levy, S. M. (1998). Medical conditions associated with missing teeth and edentulism in the institutionalized elderly. *Special Care in Dentistry*, 18(3), 123-127.

57. Papas, A. S., Joshi, A., Giunta, J. L., & Palmer, C. A. (1998). Relationships among education, dentate status, and diet in adults. *Special Care in Dentistry*, 18(1), 26-32.
58. Hamada, M. O., Garrett, N. R., Roumanas, E. D., Kapur, K. K., Freymiller, E., Han, T., ... & Levin, S. (2001). A randomized clinical trial comparing the efficacy of mandibular implant-supported overdentures and conventional dentures in diabetic patients. Part IV: Comparisons of dietary intake. *The Journal of prosthetic dentistry*, 85(1), 53-60.
59. Kapur, K. K., & Okubo, J. (1970). Effect of impaired mastication on the health of rats. *Journal of dental research*, 49(1), 61-68.
60. Levine, A. S., & Silvis, S. E. (1980). Absorption of whole peanuts, peanut oil, and peanut butter. *New England Journal of Medicine*, 303(16), 917-918.
61. Hoebler, C., Karinithi, A., Devaux, M. F., Guillon, F., Gallant, D. J. G., Bouchet, B., ... & Barry, J. L. (1998). Physical and chemical transformations of cereal food during oral digestion in human subjects. *British Journal of Nutrition*, 80(05), 429-436.
62. Rodriguez-Oliveros, A. (1947). Gastritis in the toothless. *The Review of gastroenterology*, 14(3), 180.
63. Hedde, R. D., Lindsey, T. O., Parish, R. C., Daniels, H. D., Morgenthien, E. A., & Lewis, H. B. (1985). Effect of diet particle size and feeding of H2-receptor antagonists on gastric ulcers in swine. *Journal of animal science*, 61(1), 179-186.
64. Chen, M. K., & Lowenstein, F. (1984). Masticatory handicap, socioeconomic status, and chronic conditions among adults. *The Journal of the American Dental Association*, 109(6), 916-918.
65. Mattila, K. J., Nieminen, M. S., Valtonen, V. V., Rasi, V. P., Kesäniemi, Y. A., Syrjälä, S. L., ... & Jokinen, M. J. (1989). Association between dental health and acute myocardial infarction. *Bmj*, 298(6676), 779-781.
66. DeStefano, F., Anda, R. F., Kahn, H. S., Williamson, D. F., & Russell, C. M. (1993). Dental disease and risk of coronary heart disease and mortality. *Bmj*, 306(6879), 688-691.
67. Beck, J., Garcia, R., Heiss, G., Vokonas, P. S., & Offenbacher, S. (1996). Periodontal disease and cardiovascular disease. *Journal of periodontology*, 67(10s), 1123-1137.
68. Joshipura, K. J., Rimm, E. B., Douglass, C. W., Trichopoulos, D., Ascherio, A., & Willett, W. C. (1996). Poor oral health and coronary heart disease. *Journal of dental research*, 75(9), 1631-1636.
69. Joshipura, K. J., Douglass, C. W., & Willett, W. C. (1998). Possible explanations for the tooth loss and cardiovascular disease relationship. *Annals of Periodontology*, 3(1), 175-183.
70. Correa, P., Haenszel, W., Cuello, C., Tannenbaum, S., & Archer, M. (1975). A model for gastric cancer epidemiology. *The Lancet*, 306(7924), 58-60.
71. Negri, E., La Vecchia, C., Franceschi, S., D'Avanzo, B., & Parazzini, F. (1991). Vegetable and fruit consumption and cancer risk. *International journal of cancer*, 48(3), 350-354.
72. Steinmetz, K. A., & Potter, J. D. (1991). Vegetables, fruit, and cancer. II. Mechanisms. *Cancer Causes & Control*, 2(6), 427-442.
73. Steinmetz, K. A., & Potter, J. D. (1991). Vegetables, fruit, and cancer. I. Epidemiology. *Cancer Causes & Control*, 2(5), 325-357.
74. Block, G., Patterson, B., & Subar, A. (1992). Fruit, vegetables, and cancer prevention: a review of the epidemiological evidence. *Nutrition and cancer*, 18(1), 1-29.
75. Willett, W. C. (1994). Diet and health: what should we eat?. *Science*, 264(5158), 532-537.
76. Tavani, A., & La Vecchia, C. (1995). Fruit and vegetable consumption and cancer risk in a Mediterranean population. *The American journal of clinical nutrition*, 61(6), 1374S-1377S.
77. Scheppach, W., Bingham, S., Boutron-Ruault, M. C., de Verdier, M. G., Moreno, V., Nagengast, F. M., ... & Kasper, H. (1999). WHO Consensus statement on the role of nutrition in colorectal cancer*. *European Journal of Cancer Prevention*, 8(1), 57-62.
78. Jansen, M. C., Bueno-de-Mesquita, H. B., Buzina, R., Fidanza, F., Menotti, A., Blackburn, H., ... & Kromhout, D. (1999). Dietary fiber and plant foods in relation to colorectal cancer mortality: the Seven Countries Study. *International journal of cancer*, 81(2), 174-179.
79. Schneeman, B. O. (1987). Dietary fiber and gastrointestinal function. *Nutrition reviews*, 45(7), 129-132.
80. Jenkins, D. J., Jenkins, A. L., Wolever, T. M., Rao, A. V., & Thompson, L. U. (1986). Fiber and starchy foods: gut function and implications in disease. *The American journal of gastroenterology*, 81(10), 920-930.
81. Werler, M. M., Shapiro, S., & Mitchell, A. A. (1993). Periconceptional folic acid exposure and risk of occurrent neural tube defects. *Jama*, 269(10), 1257-1261.
82. Steinberg, D., & Witztum, J. L. (1990). Lipoproteins and atherogenesis: current concepts. *Jama*, 264(23), 3047-3052.
83. Ulbricht, T. L. V., & Southgate, D. A. T. (1991). Coronary heart disease: seven dietary factors. *The Lancet*, 338(8773), 985-992.

84. Stampfer, M. J., Sacks, F. M., Salvini, S., Willett, W. C., & Hennekens, C. H. (1991). A prospective study of cholesterol, apolipoproteins, and the risk of myocardial infarction. *New England Journal of Medicine*, 325(6), 373-381.
85. Sjöström, C. D., Lissner, L., Wedel, H., & Sjöström, L. (1999). Reduction in incidence of diabetes, hypertension and lipid disturbances after intentional weight loss induced by bariatric surgery: the SOS Intervention Study. *Obesity research*, 7(5), 477-484.
86. Gumbiner, B. (1999). The treatment of obesity in type 2 diabetes mellitus. *Primary Care: Clinics in Office Practice*, 26(4), 869-883.