

A Study to Find the Efficacy of Pregabalin in Diabetic Neuropathy Pain

Dr. Shreedhar Holla Nittor

Associate Professor, Department of Internal Medicine, Srinivas Institute of Medical Sciences, Mukka, Surathkal, Mangalore, Karnataka 574146, India

DOI: [10.36348/sjm.2019.v04i08.015](https://doi.org/10.36348/sjm.2019.v04i08.015)

Received: 03.08.2019 | Accepted: 16.08.2019 | Published: 30.08.2019

*Corresponding author: Dr. Shreedhar Holla Nittor

Abstract

Diabetes is a metabolic disorder and long standing diabetic patients are prone to get neuropathy. The pain in the neuropathy is unbearable and cannot be treated by cox inhibitors. Pregabalin is a structural similar drug of the inhibitory neurotransmitter γ -amino butyric acid but it is not functionally the same. It binds to the α -2- δ subunit of voltage-gated calcium channels reducing the release of several excitatory neurotransmitters and blocking the development of hyperalgesia and central sensitization. This study puts in a effort to find the efficacy of this drug in controlling the diabetic neuropathy pain.

Keywords: Pregabalin, efficacy, Diabetes, analgesia, pain.

Copyright © 2019: This is an open-access article distributed under the terms of the Creative Commons Attribution license which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use (Non-Commercial, or CC-BY-NC) provided the original author and source are credited.

INTRODUCTION

Diabetes is a metabolic disorder and long standing diabetic patients are prone to get neuropathy. The pain in the neuropathy is unbearable and cannot be treated by cox inhibitors. Pregabalin is a structural similar drug of the inhibitory neurotransmitter γ -amino butyric acid but it is not functionally the same. It binds to the α -2- δ subunit of voltage-gated calcium channels reducing the release of several excitatory neurotransmitters and blocking the development of hyperalgesia and central sensitization.

Opioids deals with acute pain and tricyclic antidepressants (TCAs) were used for chronic neuropathic conditions [1]. Pregabalin binds to the α -2- δ subunit of voltage-gated calcium channels reducing the release of several excitatory neurotransmitters and blocking the development of hyperalgesia and central sensitization [2]. Pregabalin has anticonvulsant anti-hyperalgesic and anxiolytic properties similar to gabapentin, but it has a more favorable pharmacokinetic profile including dose-independent absorption [3]. Recently the role as oral pre-emptive analgesic of pregabalin for postoperative pain relief has been reviewed [4, 5]. This study puts in a effort to find the efficacy of this drug in controlling the diabetic neuropathy pain.

AIMS AND OBJECTIVES

To study the efficacy of pregabalin in diabetic neuropathy.

MATERIALS AND METHODS

This study was done in the Department of Internal Medicine, Srinivas Institute of Medical Sciences.

Institution Ethical Clearance was obtained

This study was done using 60 patients. All were severely in pain according to pain numeric scale. They were given adjusted dose of pregabalin for 3 weeks and then they were checked for pain scores again and then reported. The study was done from July 2017 to June 2018.

Inclusion Criteria

- The patients were aged between 30-50 years

Exclusion Criteria

- Not consented patients

Numeric Pain Scale was used:

0	No Pain
1-3	Mild Pain (nagging, annoying, interfering little with ADLs)
4-6	Moderate Pain (interferes significantly with ADLs)
7-10	Severe Pain (disabling; unable to perform ADLs)

All the statistics were done using the SPSS software 2015 (California)

RESULTS

Table-1: Age Distribution

Number	Mean age	Std Deviation
60	37.27 years	11.48 years

Table-2: Sex Distribution

Number	Male	Female
60	49	11

Table-3: Co-Morbidities

Co-Morbidities:	Frequency
HTN	07
DM	01
DM and HTN	01
Vertigo	03
Any other CNS related illness	06
Any other CVS related illness	01
Any other Respiratory illness	01
Muscle Related Illness	01

Table-4: Pain Scale after 3 weeks of adjusted dose use

	Male	Female
Mild	41	9
Moderate	07	1
Severe	01	Nil

DISCUSSION

Pregabalin has anticonvulsant anti-hyperalgesic and anxiolytic properties similar to gabapentin, but it has a more favorable pharmacokinetic profile including dose-independent absorption³. Recently the role as oral pre-emptive analgesic of pregabalin for postoperative pain relief has been reviewed [4].

It is advantageous due to high efficacy with less drug doses and less chances of aspiration pneumonitis. Due to its limitations in the form of lesser control of block height & limited duration of analgesia researchers have used battery of drugs intrathecally like vasoconstrictors, (epinephrine) opioids, (fentanyl, buprenorphine) benzodiazepines, (midazolam) ketamine and many others as adjuvant to local anaesthetics to prolong the duration of sensory block & achieve longer perioperative analgesia [6, 7]. But each of this adjuvant has certain limitations of their own hence search for better options for acute postoperative analgesia research is still continuing. Provision of effective pain relief is a prerequisite accelerated convalescence. Previously the drugs used for acute & chronic pain were categorically different. Opioids, NSAIDS & local anaesthetics were tools for dealing with acute pain and tricyclic antidepressants (TCAs) were used for chronic neuropathic conditions.

Two comparative studies gabapentin v/s pregabalin using single oral pre-emptive drug for infraumbilical surgeries under SAB, for evaluation of their comparative efficacy in terms of a cute postoperative analgesic benefits with rescue analgesic as diclofenac have shown similar results to our study [8, 9].

CONCLUSION

Pregabalin is a good drug to control diabetic neuropathy pain.

REFERENCES

- Shneker, B. F., & McAuley, J. W. (2005). Pregabalin: a new neuromodulator with broad therapeutic indications. *Annals of Pharmacotherapy*, 39(12), 2029-2037.
- Chizh, B. A., Göhring, M., Tröster, A., Quartey, G. K., Schmelz, M., & Koppert, W. (2007). Effects of oral pregabalin and aprepitant on pain and central sensitization in the electrical hyperalgesia model in human volunteers. *British journal of anaesthesia*, 98(2), 246-254.
- Guay, D. R. (2005). Pregabalin in neuropathic pain: a more "pharmaceutically elegant" gabapentin?. *The American journal of geriatric pharmacotherapy*, 3(4), 274-287.
- Frampton, J. E., & Foster, R. H. (2005). Pregabalin: In the treatment of post herpetic neuralgia. *Drugs*, 65, 111-118.
- Gilron, I. (2007). Gabapentin and pregabalin for chronic neuropathic and early postsurgical pain: current evidence and future directions. *Current Opinion in Anesthesiology*, 20(5), 456-472.
- Sahu, S., Sachan, S., Verma, A., & Pandey, H. D. (2010). Evaluation of pregabalin for attenuation of postoperative pain in below umbilical surgeries under spinal anaesthesia. *Journal of Anaesthesiology Clinical Pharmacology*, 26(2), 167-171.
- Saraswat, V., & Arora, V. (2008). Preemptive gabapentin vs pregabalin for acute postoperative pain after surgery under spinal anaesthesia. *Indian journal of anaesthesia*, 52(6), 829-834.
- Mathiesen, O., Jacobsen, L. S., Holm, H. E., Randall, S., Adamiec-Malmstroem, L., Graungaard, B. K., ... & Dahl, J. B. (2008). Pregabalin and dexamethasone for postoperative pain control: a randomized controlled study in hip arthroplasty. *British journal of anaesthesia*, 101(4), 535-541.
- Tiippana EM, Hamunen K, Kontinen VK, Kalso E. Do surgical patients benefit from perioperative gabapentin/pregabalin? A systematic review of efficacy and safety. *Anesthesia & Analgesia*. 2007 Jun 1;104(6):1545-56.