

Forestier Syndrome Presenting with Dysphagia

Brahim El Jebbouri^{1*}, Ahmed Rouihi¹, Mohamed Moutaoukel¹

¹Mohamed VI Military Hospital, Dakhla Morocco

DOI: [10.36348/sjm.2022.v07i07.008](https://doi.org/10.36348/sjm.2022.v07i07.008)

Received: 13.06.2022 | Accepted: 04.07.2022 | Published: 30.07.2022

*Corresponding Author: Brahim El Jebbouri
Mohamed VI Military Hospital, Dakhla Morocco

Abstract

Diffuse idiopathic skeletal hyperostosis (DISH) (Forestier syndrome) is a rheumatologic disease, the etiology of which is not exactly known. It is characterized by spinal osteophyte formations resulting from the ossification of the paravertebral ligaments and muscles. Anterior longitudinal ligament is the usual site of involvement and the frequency of the disease increases after the 5th decade. Lower cervical segments are the most frequently involved regions whereas the upper cervical involvement leading to dysphagia is very rare. In this report, a 65-year-old patient with Forestier syndrome in lower cervical region presenting with dysphagia was presented. Anterior cervical osteophyte resection was performed with no need for discectomy, fusion or stabilization. The patient showed a significant improvement in his all preoperative symptoms, and no recurrence was detected at 1-year follow-up.

Keywords: Forestier syndrome, DISH, rheumatologic disease, discectomy.

Copyright © 2022 The Author(s): This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

INTRODUCTION

Diffuse idiopathic skeletal hyperostosis (DISH) is a rheumatologic disease which can cause cervical osteophytes and lead to otolaryngological symptoms such as difficulty in swallowing, difficulty in breathing and loss of weight. Primarily involved structure is the anterior longitudinal ligament, ossification and calcification of which result in osteophyte formation [1]. Clinical and radiological characteristics of the disease was defined by Forestier and Rotés-Querol for the first time in 1950 [2], and the disease was named as DISH in 1975, by Resnick *et al*, [3]. In DISH, cervical area is less affected compared to thoracic and lumbar vertebral segments. The most commonly affected area is C4–7 (i.e., lower cervical) segments which correspond to relatively more mobile regions of cervical vertebra [4]. However, Forestier syndrome is very presenting with dysphagia [5, 6]. Here we report a very rare case of Forestier syndrome with lower cervical involvement presenting with dysphagia as a result of osteophyte at C4–5 level. The patient underwent anterior cervical osteophyte resection with no discectomy, fusion or stabilization.

CASE PRESENTATION

A 65-year-old male patient was referred to our department with the complaints of difficulty in

swallowing, difficulty in breathing and severe loss of weight (10 kg in 6 months). Medical history and physical examination were unremarkable. There was no neurological involvement. Laboratory tests including HLA-B27 were within normal limits. Extensive ossification of the anterior longitudinal ligament starting from anterior middle C4 extending to anterior C5 vertebrae. Hyperostosis with osteophytic formation was especially apparent at C4–5 level. In the cervical lateral X-Ray (Figure1). Due to severe symptoms such as difficulty in breathing, dysphagia and excessive loss of weight the patient was operated on. Cervical osteophyte resection was performed between C4–5 via an anterolateral approach. No postoperative complication was encountered. Respiratory difficulty and difficulty in taking fluids resolved immediately after the surgery. Difficulty in swallowing solid foods declined but still existed in the early postoperative period. Postoperative cervical X-Ray revealed that osteophytic formations had been sufficiently removed (Figure 2). The patient was discharged on the postoperative 4th day without any neurological deficit. Difficulty in swallowing solid foods gradually improved and was completely resolved at the end of the third month of operation. The patient was free of symptoms at 1-year follow-up examination.



FIG- 1



FIG- 2

DISCUSSION

DISH is a rare pathology characterized by ossification and calcification of paravertebral ligaments and muscles [3]. DISH is observed more frequently in males than females and its frequency increases with age [3, 7]. It has a prevalence of 8–10% over the age of 65 and, it is rarely seen under the age of 45 [3]. Etiology of the disease is not known exactly, however, it has been reported that DISH is associated with acromegaly, type II diabetes mellitus, obesity, genetic factors such as hypervitaminosis A, HLA-B27, HLA-B5, HLA-A11, ankylosing spondylitis and infectious diseases [1, 5]. The medical history of our case was unremarkable, no associated disease was detected and he was HLA-B27 negative. Spine, pelvis, patella and calcaneus are the most frequently involved regions in DISH [8]. Sacroiliac joint involvement is not observed in DISH [3]. Throughout the spine; it is observed 97% in thoracic, 90% in lumbar, 78% in cervical vertebrae, and 70% in these three regions together [9]. Anterior longitudinal ligament is the usual site of involvement in vertebral column. Due to immobility of the thoracic vertebrae, osteophytic structures extend continuously along the thoracic segments without any interruption. But in the cervical and lumbar areas, hyperostosis is rather horizontal [3]. The most frequently involved region in the cervical vertebrae is the region between C4–7. On the other hand, involvement of the upper cervical vertebrae is very rare in DISH [10]. In our case, the site of involvement was C2–5, which is very rarely reported in the literature. The osteophyte located anterior to vertebra corpus, at the level of C2–3, was the main cause of dysphagia. The most frequently observed symptoms of DISH are neurovascular compression, restriction of movement, cervicalgia, stridor, hoarseness, and dysphagia [11-13]. Dysphagia is observed in 0.6–1.0% of DISH cases, where cervical area is the primarily affected region (14). Dysphagia typically appears as a result of esophageal compression in lower cervical segments, most frequently at C4–5 level [10]. Esophageal occlusion, recurrent laryngeal nerve damage, esophageal fibrosis, and inflammation have been considered among the reasons of dysphagia [15]. In the case we are presenting, osteophytic structures were determined in anterior cervical area, between C2–5. Our case, which represents an upper cervical involvement, is a rare case containing C2–3 osteophyte leading to dysphagia. DISH is diagnosed based on radiological examinations [12, 13]. There are three radiological diagnostic criteria: (I) formation of calcification and bone bridges in two adjacent vertebrae; (II) intervertebral disk height being normal or slightly reduced; (III) absence of ankylosis in the interapophysial joints [3, 14]. In lateral X-ray, hyperostosis is observed only on the anterior surface of the vertebra in the beginning of the disease, while in the latter stages; osteophytes are firmly adhered to the cortex, starting from anterior border to the intervertebral area [11]. CT is the gold standard diagnostic tool [4]. Especially sagittal, coronal and three-dimensional

reconstructions of CT allow us to clearly identify the relation of osteophytes with the surrounding anatomical structures. Barium swallowing study is a useful method in determining the degree and the localization of esophageal compression [7]. In our case; lateral X-ray, CT and barium swallowing studies were performed to make the diagnosis. In symptomatic DISH cases, conservative treatment options are recommended if clinical findings are not very severe [11]. Non-steroid anti-inflammatory drugs, steroids, myorelaxants, dietary changes, anti-reflux treatments are among the conservative treatment options [1, 8]. Surgical treatment should be considered in cases with severe symptoms such as progressive dysphagia, excessive loss of weight, difficulty in breathing and in patients who do not respond to the conservative treatment [12]. In our case, surgery was indicated due to gradually progressing severe dysphagia, loss of weight and respiratory distress. It is important to distinguish hyperostotic structures from osteoarthritic osteophytes since each has a different surgical treatment. Removal of hyperostotic structures is sufficient in surgical treatment of DISH; while additional stabilization of vertebra is required after removal of osteoarthritic osteophytes [3, 9]. Anterior cervical approach, posterolateral approach and transpharyngeal approaches are the surgical methods that can be preferred. Anterior cervical approach is considered safe and effective [10]; however, it may lead to some complications such as recurrent laryngeal nerve injury, Horner syndrome, fistula formation and instability [1]. Although posterolateral approach provides a wide surgical exposure, it has the risk of injury to internal carotid artery and sympathetic chain due to excessive retraction [14]. Transpharyngeal approach is another surgical approach that can be preferred [10]. The most important complication of this method is the infections that occur as a result of the contamination by the oropharyngeal flora. In our case, osteophytic structures were removed via an anterior cervical approach. It has been seen that sufficient exposure can be obtained between C2–5 with an anterior cervical approach. The prognosis varies in the literature. Hirano *et al.*, [9] reported 65% recurrence rate after a mean postoperative follow-up period of 4.5 years. Miyamoto *et al.*, [15] reported seven patients with a mean postoperative follow-up period of 9 years and detected 100% recurrence of osteophytes, with a rate of approximately 1 mm/year. Recurrence was not detected during 1-year follow-up period of our case.

CONCLUSIONS

Forestier syndrome presenting with dysphagia is very rare. Surgery is indicated when the symptoms are severe and progressive. Despite the possibility of recurrence in long-term follow-up, resection of osteophytes via an anterior cervical approach is a reliable and effective surgical treatment modality in Forestier syndrome.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

REFERENCES

1. Akhtar, S., O'flynn, P. E., Kelly, A., & Valentine, P. M. W. (2000). The management of dysphasia in skeletal hyperostosis. *The Journal of Laryngology & Otolaryngology*, 114(2), 154-157.
2. Forestier, J., & Rotés-Querol, J. (1950). Senile ankylosing hyperostosis of the spine. *Annals of the rheumatic diseases*, 9(4), 321-330.
3. Resnick, D., Shaul, S. R., & Robins, J. M. (1975). Diffuse idiopathic skeletal hyperostosis (DISH): Forestier's disease with extraspinal manifestations. *Radiology*, 115(3), 513-524.
4. Özkırış, M., Okur, A., Kapusuz, Z., & Saydam, L. (2014). Forestier's syndrome: a rare cause of dysphagia. *Kulak Burun Bogaz İhtisas Dergisi: KBB= Journal of Ear, Nose, and Throat*, 24(1), 54-57.
5. Mader, R. (2008). Diffuse idiopathic skeletal hyperostosis: time for a change. *The Journal of Rheumatology*, 35(3), 377-379.
6. Solaroğlu, I., Okutan, O., Karakuş, M., SAYGILI, B., & Beşkonaklı, E. (2008). Dysphagia due to diffuse idiopathic skeletal hyperostosis of the cervical spine. *Turkish Neurosurgery*, 18(4), 409-411.
7. Kiss, C., O'Neill, T. W., Mitzszova, M., Szilágyi, M., Donath, J., & Poór, G. Y. (2002). Prevalence of diffuse idiopathic skeletal hyperostosis in Budapest, Hungary. *Rheumatology*, 41(11), 1335-1336.
8. Andersen, P. M., & Fagerlund, M. (2000). Vertebrogenic dysphagia and gait disturbance mimicking motor neuron disease. *Journal of Neurology, Neurosurgery & Psychiatry*, 69(4), 560-561.
9. Hirano, H., Suzuki, H., Sakakibara, T., Higuchi, Y., Inoue, K., & Suzuki, Y. (1982). Dysphagia due to hypertrophic cervical osteophytes. *Clinical Orthopaedics and Related Research*, 167, 168-172.
10. Carrau, R. L., Cintron, F. R., & Astor, F. (1990). Transcervical approaches to the prevertebral space. *Archives of Otolaryngology-Head & Neck Surgery*, 116(9), 1070-1073.
11. Ebo, D., Goethals, L., Bracke, P., Mercelis, R., & De Clerck, L. S. (2000). Dysphagia in a patient with giant osteophytes: case presentation and review of the literature. *Clinical Rheumatology*, 19(1), 70-72.
12. Hwang, J. S., Chough, C. K., & Joo, W. I. (2013). Giant anterior cervical osteophyte leading to Dysphagia. *Korean Journal of Spine*, 10(3), 200-202.
13. Kmucha, S. T., & Cravens Jr, R. B. (1994). DISH syndrome and its role in

- dysphagia. *Otolaryngology—Head and Neck Surgery*, 110(4), 431-436.
14. McCafferty, R. R., Harrison, M. J., Tamas, L. B., & Larkins, M. V. (1995). Ossification of the anterior longitudinal ligament and Forestier's disease: an analysis of seven cases. *Journal of neurosurgery*, 83(1), 13-17.
 15. Miyamoto, K., Sugiyama, S., Hosoe, H., Iinuma, N., Suzuki, Y., & Shimizu, K. (2009). Postsurgical recurrence of osteophytes causing dysphagia in patients with diffuse idiopathic skeletal hyperostosis. *European Spine Journal*, 18(11), 1652-1658.